


AMACR Antibody (V2506)


Catalog No.	Formulation	Size
V2506-100UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide	100 ug
V2506-20UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide	20 ug
V2506SAF-100UG	1 mg/ml in 1X PBS; BSA free, sodium azide free	100 ug
V2506IHC-7ML	Prediluted in 1X PBS, 0.05% sodium azide; *For IHC use only*	7 ml

[Bulk quote request](#)


Availability	1-3 business days
Species Reactivity	Human
Format	Purified
Clonality	Polyclonal (rabbit origin)
Isotype	Rabbit IgG
Purity	Protein A affinity chromatography
UniProt	Q9UHK6
Localization	Cytoplasmic
Applications	Immunohistochemistry (FFPE) : 1:50-1:100 for 30-60 min at RT Western Blot : 1:50-1:100
Limitations	This AMACR antibody is available for research use only.



Formalin-fixed, paraffin-embedded human prostate carcinoma stained with AMACR antibody.



Formalin-fixed, paraffin-embedded human lung carcinoma stained with AMACR antibody.



Western blot testing of human kidney lysate with AMACR antibody. Predicted molecular weight ~43 kDa.

Description

This antibody recognizes a protein of 44kDa, which is identified as AMACR, also known as p504S. It is an enzyme that is involved in bile acid biosynthesis and beta-oxidation of branched-chain fatty acids. AMACR is essential in lipid metabolism. It is expressed in cells of premalignant high-grade prostatic intraepithelial neoplasia (HGPIN) and prostate adenocarcinoma. The majority of the carcinoma cells show a distinct granular cytoplasmic staining reaction. AMACR is present at low or undetectable levels in glandular epithelial cells of normal prostate and benign prostatic hyperplasia. A spotty granular cytoplasmic staining is seen in a few cells of the benign glands. AMACR is expressed in normal liver (hepatocytes), kidney (tubular epithelial cells) and gall bladder (epithelial cells). Expression has also been found in lung (bronchial epithelial cells) and colon (colonic surface epithelium). AMACR expression can also be found in hepatocellular carcinoma and kidney carcinoma. Past studies have also shown that AMACR is expressed in various colon carcinomas (well, moderately and poorly differentiated) and over expressed in prostate carcinoma.

Application Notes

Optimal dilution of the AMACR antibody should be determined by the researcher.

1. Staining of formalin-fixed tissues requires boiling tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 10-20 min followed by cooling at RT for 20 minutes
2. The prediluted format is supplied in a dropper bottle and is optimized for use in IHC. After epitope retrieval step (if required), drip mAb solution onto the tissue section and incubate at RT for 30 min.

Immunogen

A synthetic peptide from the human protein was used as the immunogen for the AMACR antibody.

Storage

Store the AMACR antibody at 2-8°C (with azide) or aliquot and store at -20°C or colder (without azide).

