


Anti-Progesterone Receptor Antibody [clone SPM566] (V9066)

Catalog No.	Formulation	Size
V9066-100UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide	100 ug
V9066-20UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide	20 ug
V9066SAF-100UG	1 mg/ml in 1X PBS; BSA free, sodium azide free	100 ug
V9066IHC-7ML	Prediluted in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide; *For IHC use only*	7 ml

[Bulk quote request](#)

Availability	1-3 business days
Species Reactivity	Human
Format	Purified
Host	Mouse
Clonality	Monoclonal (mouse origin)
Isotype	Mouse IgG1, kappa
Clone Name	SPM566
Purity	Protein G affinity chromatography
UniProt	P06401
Localization	Nuclear, cytoplasmic
Applications	Immunohistochemistry (FFPE) : 1-2ug/ml for 30 min at RT
Limitations	This anti-Progesterone Receptor antibody is available for research use only.



IHC: Formalin-fixed, paraffin-embedded human breast carcinoma stained with anti-Progesterone Receptor antibody (clone SPM566).

Description

This mAb is specific to progesterone receptor and shows minimal cross-reaction with other members of the family. Progesterone receptor is expressed as two major isoforms, PR-A (81kDa) and PR-B (116kDa). Expression of PgR has been suggested to reflect an intact estrogen regulatory machinery and therefore, predict better clinical response to endocrine therapy than ER alone.

Application Notes

The optimal dilution of the anti-Progesterone Receptor antibody for each application should be determined by the researcher.

1. Staining of formalin-fixed tissues requires boiling tissue sections in pH 9 10mM Tris with 1mM EDTA for 10-20 min followed by cooling at RT for 20 minutes.
2. The prediluted format is supplied in a dropper bottle and is optimized for use in IHC. After epitope retrieval step (if required), drip mAb solution onto the tissue section and incubate at RT for 30 min.

Immunogen

Recombinant human protein was used as the immunogen for this anti-Progesterone Receptor antibody.

Storage

Store the anti-Progesterone Receptor antibody at 2-8oC (with azide) or aliquot and store at -20oC or colder (without azide).