


## APEX2 Antibody (F55013)


Catalog No.	Formulation	Size
F55013-0.4ML	In 1X PBS, pH 7.4, with 0.09% sodium azide	0.4 ml
F55013-0.08ML	In 1X PBS, pH 7.4, with 0.09% sodium azide	0.08 ml

[Bulk quote request](#)


<b>Availability</b>	1-3 business days
<b>Species Reactivity</b>	Human
<b>Format</b>	Purified
<b>Host</b>	Rabbit
<b>Clonality</b>	Polyclonal (rabbit origin)
<b>Isotype</b>	Rabbit Ig
<b>Purity</b>	Antigen affinity purified
<b>UniProt</b>	Q9UBZ4
<b>Localization</b>	Cytoplasmic, nuclear
<b>Applications</b>	Immunofluorescence : 1:25 Flow Cytometry : 1:10-1:50 (1x10e6 cells) Immunohistochemistry (FFPE) : 1:10-1:50 Western Blot : 1:500-1:1000
<b>Limitations</b>	This APEX2 antibody is available for research use only.




IHC testing of FFPE human lung carcinoma tissue with APEX2 antibody. HIER: steam section in pH6 citrate buffer for 20 min and allow to cool prior to staining.



Immunofluorescent staining of human U-251 cells with APEX2 antibody (green) and anti-Actin (red).



Western blot testing of human MCF7 cell lysate with APEX2 antibody. Predicted molecular weight ~57 kDa.



Flow cytometry testing of human MCF7 cells with APEX2 antibody; Blue=isotype control, Green= APEX2 antibody.

## Description

APEX2 may participate in both nuclear and mitochondrial post-replicative base excision repair (BER). In the nucleus functions in the PCNA-dependent BER pathway.

## Application Notes

The stated application concentrations are suggested starting points. Titration of the APEX2 antibody may be required due to differences in protocols and secondary/substrate sensitivity.

## Immunogen

A portion of amino acids 143-171 from the human protein was used as the immunogen for the APEX2 antibody.

## Storage

Aliquot the APEX2 antibody and store frozen at -20oC or colder. Avoid repeated freeze-thaw cycles.