

## ATG4D Antibody (F54717)


Catalog No.	Formulation	Size
F54717-0.4ML	In 1X PBS, pH 7.4, with 0.09% sodium azide	0.4 ml
F54717-0.08ML	In 1X PBS, pH 7.4, with 0.09% sodium azide	0.08 ml

[Bulk quote request](#)


<b>Availability</b>	1-3 business days
<b>Species Reactivity</b>	Human, Mouse
<b>Format</b>	Purified
<b>Clonality</b>	Polyclonal (rabbit origin)
<b>Isotype</b>	Rabbit Ig
<b>Purity</b>	Purified
<b>UniProt</b>	Q86TL0
<b>Localization</b>	Cytoplasmic
<b>Applications</b>	Western Blot : 1:500-1:2000 Immunohistochemistry (FFPE) : 1:25
<b>Limitations</b>	This ATG4D antibody is available for research use only.




Western blot testing of human HL60 cell lysate with ATG4D antibody. Expected molecular weight: 53-56 kDa.



Western blot testing of mouse kidney tissue lysate with ATG4D antibody. Expected molecular weight: 53-56 kDa.



Western blot testing of 1) non-transfected and 2) transfected 293 cell lysate with ATG4D antibody.



IHC testing of FFPE human cancer tissue with ATG4D antibody. HIER: steam section in pH6 citrate buffer for 20 min and allow to cool prior to staining.

## Description

Macroautophagy is the major inducible pathway for the general turnover of cytoplasmic constituents in eukaryotic cells, it is also responsible for the degradation of active cytoplasmic enzymes and organelles during nutrient starvation.

Macroautophagy involves the formation of double-membrane bound autophagosomes which enclose the cytoplasmic constituent targeted for degradation in a membrane bound structure, which then fuse with the lysosome (or vacuole) releasing a single-membrane bound autophagic bodies which are then degraded within the lysosome (or vacuole). APG4 is a cysteine protease required for autophagy, which cleaves the C-terminal part of either MAP1LC3, GABARAPL2 or GABARAP, allowing the liberation of form I. A subpopulation of form I is subsequently converted to a smaller form (form II). Form II, with a revealed C-terminal glycine, is considered to be the phosphatidylethanolamine (PE)-conjugated form, and has the capacity for the binding to autophagosomes.

## Application Notes

The stated application concentrations are suggested starting points. Titration of the ATG4D antibody may be required due to differences in protocols and secondary/substrate sensitivity.

## Immunogen

A portion of amino acids 220-249 from the human protein was used as the immunogen for the ATG4D antibody.

## Storage

Aliquot the ATG4D antibody and store frozen at -20oC or colder. Avoid repeated freeze-thaw cycles.