


## AURKC Antibody (F53762)


Catalog No.	Formulation	Size
F53762-0.4ML	In 1X PBS, pH 7.4, with 0.09% sodium azide	0.4 ml
F53762-0.08ML	In 1X PBS, pH 7.4, with 0.09% sodium azide	0.08 ml

[Bulk quote request](#)

<b>Availability</b>	1-3 business days
<b>Species Reactivity</b>	Human
<b>Format</b>	Purified
<b>Clonality</b>	Polyclonal (rabbit origin)
<b>Isotype</b>	Rabbit Ig
<b>Purity</b>	Purified
<b>UniProt</b>	Q9UQB9
<b>Applications</b>	Western Blot : 1:1000 IHC (Paraffin) : 1:50-1:100
<b>Limitations</b>	This AURKC antibody is available for research use only.



Western blot testing of AURKC antibody and A375 cell lysate. Predicted molecular weight ~ 33 kDa.



IHC analysis of FFPE human breast carcinoma tissue stained with the AURKC antibody

## Description

Chromosomal segregation during mitosis as well as meiosis is regulated by kinases and phosphatases. The Aurora kinases, members of the Ser/Thr protein kinase family, associate with microtubules during chromosome movement and segregation. Aurora kinase C may play a part in organizing microtubules in relation to the function of the centrosome/spindle pole during mitosis. This protein is localized to centrosome from anaphase to cytokinesis. Expression is limited to testis in normal cells. Elevated expression levels are seen only in a subset of cancer cells such as HepG2, HuH7 and HeLa cells. Aurora-C expression is maximum at M phase.

## Application Notes

Titration of the AURKC antibody may be required due to differences in protocols and secondary/substrate sensitivity.

## Immunogen

A portion of the C-terminal region of the human protein was used as immunogen for this AURKC antibody.

## Storage

Aliquot the AURKC antibody and store frozen at -20oC or colder. Avoid repeated freeze-thaw cycles.