

## CCL3 Antibody / MIP-1 alpha (R31933)

Catalog No.	Formulation	Size
R31933	0.5mg/ml if reconstituted with 0.2ml sterile DI water	100 ug

**Bulk quote request**

<b>Availability</b>	1-3 business days
<b>Species Reactivity</b>	Human
<b>Format</b>	Antigen affinity purified
<b>Host</b>	Rabbit
<b>Clonality</b>	Polyclonal (rabbit origin)
<b>Isotype</b>	Rabbit IgG
<b>Purity</b>	Antigen affinity
<b>Buffer</b>	Lyophilized from 1X PBS with 2.5% BSA and 0.025% sodium azide
<b>UniProt</b>	P10147
<b>Applications</b>	Western Blot : 0.1-0.5ug/ml ELISA : 0.1-0.5ug/ml (human protein tested); request BSA-free format for coating
<b>Limitations</b>	This CCL3 antibody is available for research use only.



Western blot testing of human Jurkat cell lysate with CCL3 antibody. Expected molecular weight: 11/8 kDa (precursor/mature).

## Description

Macrophage inflammatory protein-1 alpha (MIP-1 alpha), also called CCL3, LD78. The cDNA for CCL3 predicts a mature peptide of 69 amino acids with a molecular mass of 7,889 daltons. CCL3 is a member of a newly identified superfamily of small inducible proteins involved in inflammatory responses, wound healing and tumorigenesis. CCL3 is a chemokine that has pro-inflammatory and stem cell inhibitory activities in vitro. It constitutes an important second signal for mast cell degranulation in the conjunctiva in vivo and consequently for acute-phase disease.

## Application Notes

Optimal dilution of the CCL3 antibody should be determined by the researcher.

## Immunogen

Amino acids 24-92 of human CCL3 were used as the immunogen for the CCL3 antibody.

## Storage

After reconstitution, the CCL3 antibody can be stored for up to one month at 4oC. For long-term, aliquot and store at -20oC. Avoid repeated freezing and thawing.