

CD20 Antibody [clone 109-3C2] (V8375)

Catalog No.	Formulation	Size
V8375-100UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide	100 ug
V8375-20UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide	20 ug
V8375SAF-100UG	1 mg/ml in 1X PBS; BSA free, sodium azide free	100 ug

Bulk quote request

Availability	1-3 business days
Species Reactivity	Human
Format	Purified
Host	Mouse
Clonality	Monoclonal (mouse origin)
Isotype	Mouse IgG3, kappa
Clone Name	109-3C2
Purity	Protein G affinity chromatography
UniProt	P11836
Localization	Predominantly cell surface with some cytoplasmic
Applications	Functional Studies : order Ab without BSA ELISA : order Ab without BSA for coating
Limitations	This CD20 antibody is available for research use only.



Description

Recognizes a protein of 30-33kDa, which is identified as CD20 (Workshop V; Code CD20.12. Workshop IV; Code B17). It recognizes an extracellular domain of CD20. It is a non-Ig differentiation antigen of B-cells and its expression is restricted to normal and neoplastic B-cells, being absent from all other leukocytes and tissues. CD20 is expressed by pre B-cells and persists during all stages of B-cell maturation but is lost upon terminal differentiation into plasma cells. The protein passes through the membrane 4 times with both ends in cytoplasm and exposes one short and one longer loop to the external environment. CD20 is not glycosylated in resting B-cells and its cytoplasmic domains are differentially phosphorylated upon activation. It acts as calcium channel involved in B cell activation and cell cycle progression.

Application Notes

Optimal dilution of the CD20 antibody should be determined by the researcher.

Immunogen

Stimulated human leukocytes were used as the immunogen for the CD20 antibody.

Storage

Store the CD20 antibody at 2-8oC (with azide) or aliquot and store at -20oC or colder (without azide).