

## CD33 Antibody [clone WM53] (V2060)

Catalog No.	Formulation	Size
V2060-100UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide	100 ug
V2060-20UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide	20 ug
V2060SAF-100UG	1 mg/ml in 1X PBS; BSA free, sodium azide free	100 ug

 [Citations \(5\)](#)
[Bulk quote request](#)

Species Reactivity	Human
Format	Purified
Host	Mouse
Clonality	Monoclonal (mouse origin)
Isotype	Mouse IgG1, kappa
Clone Name	WM53
Purity	Protein G purified monoclonal antibody
Buffer	1X PBS, pH 7.4
Gene ID	945
Localization	Cell surface
Applications	Flow Cytometry : 1-2ug/10 <sup>6</sup> cells
Limitations	This <b>CD33 antibody</b> is available for research use only.



## Description

This antibody recognizes a 67kDa glycoprotein, which is identified as CD33 (HLDA IV; WS Code M-505). CD33 is a transmembrane protein of the sialic acid-binding immunoglobulin-like lectin (Siglec) family. It belongs to the immunoreceptor tyrosine-based inhibitory motif (ITIM)-containing molecules able of recruiting protein tyrosine phosphatases SHP-1 and SHP-2 to signal assemblies; these ITIMs are also used for ubiquitin-mediated removal of the receptor from the cell surface. CD33 is expressed on cells of myelomonocytic lineage, binds sialic acid residues in N- and O-glycans on cell surfaces, and is a therapeutic target for acute myeloid leukemia. CD33 is expressed on myeloid progenitors, monocytes, granulocytes, dendritic cells and mast cells. It is absent on platelets, lymphocytes, erythrocytes and hematopoietic stem cells.

## Application Notes

The concentration stated for each application is a general starting point. Variations in protocols, secondaries and substrates may require the CD33 antibody to be titrated up or down for optimal performance.

1. Not suitable for use in FFPE IHC testing.

## Immunogen

Human AML cells were used as the immunogen for this CD33 antibody.

## Storage

Store the CD33 antibody at 2-8oC (with azide) or aliquot and store at -20oC or colder (without azide).

## Alternate Names

gp67; My9; p67; Sialic acid-binding Ig-like lectin 3, SIGLEC3

## References (1)