

## CD5 Antibody [clone B-B8] (V2960)

Catalog No.	Formulation	Size
V2960-100UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide	100 ug
V2960-20UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide	20 ug
V2960SAF-100UG	1 mg/ml in 1X PBS; BSA free, sodium azide free	100 ug

 [Citations \(3\)](#)
[Bulk quote request](#)

<b>Availability</b>	1-3 business days
<b>Species Reactivity</b>	Human
<b>Format</b>	Purified
<b>Host</b>	Mouse
<b>Clonality</b>	Monoclonal (mouse origin)
<b>Isotype</b>	Mouse IgG1, kappa
<b>Clone Name</b>	B-B8
<b>Purity</b>	Protein G affinity chromatography
<b>UniProt</b>	P06127
<b>Localization</b>	Cell surface
<b>Applications</b>	Flow Cytometry : 0.5-1ug/10 <sup>6</sup> cells Immunofluorescence : 0.5-1ug/ml
<b>Limitations</b>	This CD5 antibody is available for research use only.



## Description

Recognizes a 67kDa transmembrane protein which is identified as CD5. The CD5 antigen is found on 95% of thymocytes and 72% of peripheral blood lymphocytes. In lymph nodes, the main reactivity is observed in T cell areas. Anti-CD5 is a pan T-cell marker that also reacts with a range of neoplastic B-cells, e.g. chronic lymphocytic leukemia/small lymphocytic lymphoma (CLL/SLL), mantle cell lymphoma, and a subset (~10%) of diffuse large B-cell lymphoma. CD5 aberrant expression is useful in making a diagnosis of mature T-cell neoplasms. Note that this mAb is not suitable for frozen tissues.

## Application Notes

Optimal dilution of the CD5 antibody should be determined by the researcher.

## Immunogen

T cells activated with PHA were used as the immunogen for the CD5 antibody.

## Storage

Store the CD5 antibody at 2-8oC (with azide) or aliquot and store at -20oC or colder (without azide).