


## CDC42EP3 Antibody (F55000)

Catalog No.	Formulation	Size
F55000-0.4ML	In 1X PBS, pH 7.4, with 0.09% sodium azide	0.4 ml
F55000-0.08ML	In 1X PBS, pH 7.4, with 0.09% sodium azide	0.08 ml

[Bulk quote request](#)

<b>Availability</b>	1-3 business days
<b>Species Reactivity</b>	Human
<b>Format</b>	Purified
<b>Host</b>	Rabbit
<b>Clonality</b>	Polyclonal (rabbit origin)
<b>Isotype</b>	Rabbit Ig
<b>Purity</b>	Antigen affinity purified
<b>UniProt</b>	Q9UKI2
<b>Localization</b>	Cytoplasmic
<b>Applications</b>	Flow Cytometry : 1:10-1:50 (1x10 <sup>6</sup> cells) Western Blot : 1:500-1:1000 Immunohistochemistry (FFPE) : 1:50-1:100
<b>Limitations</b>	This CDC42EP3 antibody is available for research use only.




IHC testing of FFPE human testis tissue with CDC42EP3 antibody. HIER: steam section in pH6 citrate buffer for 20 min and allow to cool prior to staining.

kDa  
72  
55  
43  
34  
26  
17

Western blot testing of human HEK293 cell lysate with CDC42EP3 antibody. Predicted molecular weight ~28 kDa.

kDa  
72  
55  
36  
28  
17 (-) (+)

Western blot testing of 1) non-transfected and 2) transfected 293 cell lysate with CDC42EP3 antibody.



Flow cytometry testing of human HEK293 cells with CDC42EP3 antibody; Blue=isotype control, Green= CDC42EP3 antibody.

## Description

CDC42EP3 is probably involved in the organization of the actin cytoskeleton. May act downstream of CDC42 to induce actin filament assembly leading to cell shape changes. It induces pseudopodia formation in fibroblasts.

## Application Notes

The stated application concentrations are suggested starting points. Titration of the CDC42EP3 antibody may be required due to differences in protocols and secondary/substrate sensitivity.

## Immunogen

A portion of amino acids 1-30 from the human protein was used as the immunogen for the CDC42EP3 antibody.

## Storage

Aliquot the CDC42EP3 antibody and store frozen at -20oC or colder. Avoid repeated freeze-thaw cycles.