


## Cdk4 Antibody (R30500)


Catalog No.	Formulation	Size
R30500	0.5mg/ml if reconstituted with 0.2ml sterile DI water	100 ug

**Bulk quote request**


<b>Availability</b>	1-3 business days
<b>Species Reactivity</b>	Human, Mouse, Rat
<b>Format</b>	Antigen affinity purified
<b>Host</b>	Rabbit
<b>Clonality</b>	Polyclonal (rabbit origin)
<b>Isotype</b>	Rabbit IgG
<b>Purity</b>	Antigen affinity
<b>Buffer</b>	Lyophilized from 1X PBS with 2.5% BSA and 0.025% sodium azide/thimerosal
<b>UniProt</b>	P30285
<b>Localization</b>	Nuclear, cytoplasmic
<b>Applications</b>	Western Blot : 0.5-1ug/ml Immunofluorescence : 5ug/ml Flow Cytometry : 1-3ug/million cells
<b>Limitations</b>	This Cdk4 antibody is available for research use only.



Immunofluorescent staining of FFPE human MCF7 cells with Cdk4 antibody (red) and Alpha Tubulin mAb (green). HIER: steam section in pH6 citrate buffer for 20 min.



Western blot testing of 1) human HeLa, 2) human K562, 3) rat PC-12, 4) mouse NIH 3T3 and 5) mouse RAW264.7 cell lysate with Cdk4 antibody. Predicted molecular weight ~34 kDa.



Flow cytometry testing of human MCF7 cells with Cdk4 antibody at 1ug/million cells (blocked with goat sera); Red=cells alone, Green=isotype control, Blue= Cdk4 antibody.

## Description

Cyclin-dependent kinase-4 (CDK4) is a protein-serine kinase involved in the cell cycle. Human cell division is regulated primarily at the G1-to-S or the G2-to-M boundaries within the cell cycle. The complexes formed by CDK4 and the D-type cyclins are involved in the control of cell proliferation during the G1 phase. CDK4 is inhibited by p16, also known as cyclin-dependent kinase inhibitor-2. CDK4 is mapped to 12q14. CDK4 expression and activity are required for cytokine responsiveness in T cells.

## Application Notes

The stated application concentrations are suggested starting amounts. Titration of the Cdk4 antibody may be required due to differences in protocols and secondary/substrate sensitivity.

## Immunogen

An amino acid sequence from the C-terminus of mouse Cdk4 (RISAFRALQHSYLYKEESDAE) was used as the immunogen for this Cdk4 antibody.

## Storage

After reconstitution, the Cdk4 antibody can be stored for up to one month at 4°C. For long-term, aliquot and store at -20°C. Avoid repeated freezing and thawing.