

DAAM1 Antibody (F54823)

Catalog No.	Formulation	Size
F54823-0.4ML	In 1X PBS, pH 7.4, with 0.09% sodium azide	0.4 ml
F54823-0.08ML	In 1X PBS, pH 7.4, with 0.09% sodium azide	0.08 ml

Bulk quote request

Availability	1-3 business days
Species Reactivity	Human, Hamster
Format	Purified
Clonality	Polyclonal (rabbit origin)
Isotype	Rabbit Ig
Purity	Antigen affinity purified
UniProt	Q9Y4D1
Localization	Cytoplasmic
Applications	Flow Cytometry : 1:10-1:50 (1x10 ⁶ cells) Immunohistochemistry (FFPE) : 1:50-1:100 Western Blot : 1:500-1:1000
Limitations	This DAAM1 antibody is available for research use only.




IHC testing of FFPE human skeletal muscle tissue with DAAM1 antibody. HIER: steam section in pH6 citrate buffer for 20 min and allow to cool prior to staining.

kDa
250
130
95
72
55

Western blot testing of human WiDr cell lysate with DAAM1 antibody. Predicted molecular weight ~123 kDa.

kDa
250
130
95
72
55

Western blot testing of hamster CHO cell lysate with DAAM1 antibody. Predicted molecular weight ~123 kDa.



Flow cytometry testing of human WiDr cells with DAAM1 antibody; Blue=isotype control, Green= DAAM1 antibody.

Description

Disheveled-associated activator of morphogenesis 1 binds to disheveled (Dvl) and Rho, and mediates Wnt-induced Dvl-Rho complex formation. [UniProt]

Application Notes

The stated application concentrations are suggested starting points. Titration of the DAAM1 antibody may be required due to differences in protocols and secondary/substrate sensitivity.

Immunogen

A portion of amino acids 341-370 from the human protein was used as the immunogen for the DAAM1 antibody.

Storage

Aliquot the DAAM1 antibody and store frozen at -20oC or colder. Avoid repeated freeze-thaw cycles.