


## DSCR1 Antibody (F40134)


Catalog No.	Formulation	Size
F40134-0.4ML	In 1X PBS, pH 7.4, with 0.09% sodium azide	0.4 ml
F40134-0.08ML	In 1X PBS, pH 7.4, with 0.09% sodium azide	0.08 ml

[Bulk quote request](#)

<b>Availability</b>	1-3 business days
<b>Species Reactivity</b>	Human
<b>Predicted Reactivity</b>	Bovine, Hamster, Mouse, Rat
<b>Format</b>	Purified
<b>Clonality</b>	Polyclonal (rabbit origin)
<b>Isotype</b>	Rabbit Ig
<b>Purity</b>	Purified
<b>UniProt</b>	P53805
<b>Applications</b>	Western Blot : 1:1000
<b>Limitations</b>	This DSCR1 antibody is available for research use only.



Endothelial cell lysate transferred to membrane was incubated with DSCR1 antibody at a 1:500 dilution. Data courtesy of Dr. Katherine Healey, NWCRF Institute, School of Biological Sciences, University of Wales Bangor.



The DSCR1 antibody used in western blot to detect DSCR1 in endothelial tissue lysate

## Description

DSCR1 interacts with calcineurin A and inhibits calcineurin-dependent signaling pathways, possibly affecting central nervous system development. The gene for this protein is located in the minimal candidate region for the Down syndrome phenotype, and is overexpressed in the brain of Down syndrome fetuses. Chronic overexpression of DSCR1 may lead to neurofibrillary tangles such as those associated with Alzheimer disease.

## Application Notes

Titration of the DSCR1 antibody may be required due to differences in protocols and secondary/substrate sensitivity.

## Immunogen

A portion of amino acids 222-252 from the human protein was used as the immunogen for this DSCR1 antibody.

## Storage

Aliquot the DSCR1 antibody and store frozen at -20oC or colder. Avoid repeated freeze-thaw cycles.