


## EpCAM Antibody [clone HEA125] (V7935)


Catalog No.	Formulation	Size
V7935-100UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide	100 ug
V7935-20UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide	20 ug
V7935SAF-100UG	1 mg/ml in 1X PBS; BSA free, sodium azide free	100 ug

### Bulk quote request


<b>Availability</b>	1-3 business days
<b>Species Reactivity</b>	Human
<b>Format</b>	Purified
<b>Host</b>	Mouse
<b>Clonality</b>	Monoclonal (mouse origin)
<b>Isotype</b>	Mouse IgG1, kappa
<b>Clone Name</b>	HEA125
<b>Purity</b>	Protein G affinity chromatography
<b>UniProt</b>	P16422
<b>Localization</b>	Cell surface, cytoplasmic
<b>Applications</b>	Immunofluorescence : 1-4ug/ml Immunohistochemistry (FFPE) : 1-2ug/ml Flow Cytometry : 1-2ug/million cells
<b>Limitations</b>	This EpCAM antibody is available for research use only.



Immunofluorescent staining of human MCF7 cells with EpCAM antibody (clone HEA125, green) and Reddot nuclear stain (red).



IHC staining of FFPE human breast carcinoma with EpCAM antibody (clone HEA125).  
HIER: boil tissue sections in pH 9 10mM Tris with 1mM EDTA for 10-20 min and allow to cool before testing.



Flow cytometry testing of human MCF7 cells with EpCAM antibody (clone HEA125);  
Gray=isotype control, Red= CF640-labeled EpCAM antibody.

## Description

Recognizes a 40-43kDa transmembrane epithelial glycoprotein, identified as epithelial specific antigen (ESA), or epithelial cellular adhesion molecule (Ep-CAM). It is expressed on baso-lateral cell surface in most simple epithelia and a vast majority of carcinomas with the exception of adult squamous epithelium, hepatocytes and gastric epithelial cells. Antibody to Ep-CAM has been used to distinguish adenocarcinoma from pleural mesothelioma and hepatocellular carcinoma. It is also useful in distinguishing serous carcinomas of the ovary from mesothelioma. It has been reported that this epithelial antigen plays an important role as a tumor-cell marker in lymph nodes from patients with esophageal carcinoma otherwise classified as node-negative.

## Application Notes

Optimal dilution of the EpCAM antibody should be determined by the researcher.

## Immunogen

Human colon cancer HT-29 cells were used as the immunogen for the EpCAM antibody.

## Storage

Store the EpCAM antibody at 2-8oC (with azide) or aliquot and store at -20oC or colder (without azide).