


FMO3 Antibody (R30857)

Catalog No.	Formulation	Size
R30857	0.5mg/ml if reconstituted with 0.2ml sterile DI water	100 ug

Bulk quote request

Availability	1-3 business days
Species Reactivity	Human, Mouse, Rat
Format	Antigen affinity purified
Host	Rabbit
Clonality	Polyclonal (rabbit origin)
Isotype	Rabbit IgG
Purity	Antigen affinity
Buffer	Lyophilized from 1X PBS with 2.5% BSA and 0.025% sodium azide/thimerosal
UniProt	P31513
Applications	Western Blot : 0.5-1ug/ml
Limitations	This FMO3 antibody is available for research use only.



Western blot testing of FMO3 antibody and Lane 1: rat liver; 2: rat liver.
Predicted/observed molecular weight ~60kDa.

Description

Flavin-containing Monooxygenase 3 is an enzyme that in humans is encoded by the FMO3 gene. The mammalian flavin-containing monooxygenases (FMOs) represent a multigene family whose gene products are localized in the endoplasmic reticulum of many tissues. Using quantitative RNase protection assays, FMO3 is present in low abundance in fetal liver and lung and in adult kidney and lung, and in much greater abundance in adult liver. By Western blot analysis of human liver microsomal samples ranging from 8 weeks gestation to 18 years of age, FMO1 is the major fetal isoform and FMO3 is the major adult isoform. Flavin-containing Monooxygenase 3 was expressed at intermediate levels until 11 years of age when a gender-independent increase in expression was observed during puberty. Sufferers of trimethylaminuria may

display a reduced ability to metabolize substrates for FMO3 such as nicotine, amphetamine, clozapine, deprenyl, metamphetamine, tamoxifen, ethionamide, thiacetazone, and sulindac sulfide.

Application Notes

The stated application concentrations are suggested starting amounts. Titration of the FMO3 antibody may be required due to differences in protocols and secondary/substrate sensitivity.

Immunogen

An amino acid sequence from the N-terminus of human Flavin-containing Monooxygenase 3 (DDFPNFMHNSKIQEY) was used as the immunogen for this FMO3 antibody.

Storage

After reconstitution, the FMO3 antibody can be stored for up to one month at 4oC. For long-term, aliquot and store at -20oC. Avoid repeated freezing and thawing.