


## GAP43 Antibody (RQ5884)


Catalog No.	Formulation	Size
RQ5884	0.5mg/ml if reconstituted with 0.2ml sterile DI water	100 ug

**Bulk quote request**


<b>Availability</b>	1-3 business days
<b>Species Reactivity</b>	Human, Mouse, Rat
<b>Format</b>	Antigen affinity purified
<b>Host</b>	Rabbit
<b>Clonality</b>	Polyclonal (rabbit origin)
<b>Isotype</b>	Rabbit IgG
<b>Purity</b>	Affinity purified
<b>Buffer</b>	Lyophilized from 1X PBS with 2% Trehalose and 0.025% sodium azide
<b>UniProt</b>	P06837
<b>Applications</b>	Western Blot : 0.5-1ug/ml Immunohistochemistry : 1-2ug/ml Flow Cytometry : 1-3ug/million cells Direct ELISA : 0.1-0.5ug/ml
<b>Limitations</b>	This GAP43 antibody is available for research use only.




IHC staining of FFPE rat brain with GAP43 antibody. HIER: boil tissue sections in pH8 EDTA for 20 min and allow to cool before testing.



Western blot testing of 1) rat brain, 2) mouse brain and 3) rat C6 lysate with GAP43 antibody. Expected molecular weight ~43 kDa.



Flow cytometry testing of human Raji cells with GAP43 antibody at 1ug/million cells (blocked with goat sera); Red=cells alone, Green=isotype control, Blue= GAP43 antibody.



IHC staining of FFPE human brain tissue with GAP43 antibody. HIER: boil tissue sections in pH8 EDTA for 20 min and allow to cool before testing.

## Description

Growth Associated Protein 43 (GAP43) is a protein encoded by the GAP43 gene in humans. It is mapped to 3q13.31. The protein encoded by this gene has been termed a 'growth' or 'plasticity' protein because it is expressed at high levels in neuronal growth cones during development and axonal regeneration. This protein is considered a crucial component of an effective regenerative response in the nervous system. Alternatively spliced transcript variants encoding distinct isoforms have been found for this gene.

## Application Notes

Optimal dilution of the GAP43 antibody should be determined by the researcher.

## Immunogen

Recombinant mouse protein (amino acids M1-A227) was used as the immunogen for the GAP43 antibody.

## Storage

After reconstitution, the GAP43 antibody can be stored for up to one month at 4°C. For long-term, aliquot and store at -20°C. Avoid repeated freezing and thawing.

