


## GDNF Antibody (F51281)


| Catalog No.   | Formulation                                | Size    |
|---------------|--|---------|
| F51281-0.4ML  | In 1X PBS, pH 7.4, with 0.09% sodium azide | 0.4 ml  |
| F51281-0.08ML | In 1X PBS, pH 7.4, with 0.09% sodium azide | 0.08 ml |

[Bulk quote request](#)


|                    |   |
|--------------------|---|
| Availability       | 1-3 business days   |
| Species Reactivity | Human, Mouse  |
| Format             | Antigen affinity purified   |
| Host               | Rabbit  |
| Clonality          | Polyclonal (rabbit origin)  |
| Isotype            | Rabbit Ig   |
| Purity             | Antigen affinity  |
| UniProt            | P39905  |
| Applications       | Western Blot : 1:1000<br>IHC (Paraffin) : 1:10-1:50<br>Flow Cytometry : 1:10-1:50 |
| Limitations        | This GDNF antibody is available for research use only.                            |



IHC analysis of FFPE human brain tissue stained with GDNF antibody



Western blot analysis of GDNF antibody and mouse NIH3T3 lysate.



GDNF antibody flow cytometric analysis of 293 cells (right histogram) compared to a negative control (left histogram). FITC-conjugated goat-anti-rabbit secondary Ab was used for the analysis.

## Description

GDNF is a highly conserved neurotrophic factor. The recombinant form of this protein was shown to promote the survival and differentiation of dopaminergic neurons in culture, and was able to prevent apoptosis of motor neurons induced by axotomy. This protein is processed to a mature secreted form that exists as a homodimer. The mature form of the protein is a ligand for the product of the RET (rearranged during transfection) protooncogene. In addition to the transcript encoding GDNF, two additional alternative transcripts encoding distinct proteins, referred to as astrocyte-derived trophic factors, have also been described.

## Application Notes

Titration of the GDNF antibody may be required due to differences in protocols and secondary/substrate sensitivity.

## Immunogen

A portion of amino acids 8-36 from the human protein was used as the immunogen for this GDNF antibody.

## Storage

Aliquot the GDNF antibody and store frozen at -20oC or colder. Avoid repeated freeze-thaw cycles.