


## Granzyme B Antibody (V2564)

Catalog No.	Formulation	Size
V2564-100UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide	100 ug
V2564-20UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide	20 ug
V2564SAF-100UG	1 mg/ml in 1X PBS; BSA free, sodium azide free	100 ug

[Bulk quote request](#)

<b>Availability</b>	1-3 business days
<b>Species Reactivity</b>	Human. Other species not known.
<b>Format</b>	Purified
<b>Clonality</b>	Polyclonal (rabbit origin)
<b>Isotype</b>	Rabbit IgG
<b>Purity</b>	Protein A affinity chromatography
<b>UniProt</b>	P10144
<b>Localization</b>	Cytoplasmic
<b>Applications</b>	Flow Cytometry : 0.5-1ug/million cells in 0.1ml Immunofluorescence : 0.5-1ug/ml Western Blot : 1-2ug/ml Immunohistochemistry (FFPE) : 0.5-1ug/ml for 30 min at RT (1)
<b>Limitations</b>	This Granzyme B antibody is available for research use only.



Western blot analysis of human stomach lysate using Granzyme B antibody. Predicted molecular weight: 29-37 kDa.

## Description

Granzyme B is a member of the granule serine protease family stored specifically in NK cells or cytotoxic T cells. Cytolytic

T lymphocytes (CTL) and natural killer (NK) cells share the ability to recognize, bind, and lyse specific target cells. They are thought to protect their host by lysing cells bearing on their surface 'non-self' antigens, usually peptides or proteins resulting from infection by intracellular pathogens. Granzyme B is crucial for the rapid induction of target cell apoptosis by CTLs in the cell-mediated immune response. Granzyme B is useful as a marker in the identification of NK/T-cell lymphomas. High percentages of cytotoxic T-cells have been shown to be an unfavorable prognostic indicator in Hodgkin's Disease.

## Application Notes

Optimal dilution of the Granzyme B antibody should be determined by the researcher.

1. Staining of formalin-fixed tissues requires boiling tissue sections in 10mM Citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 min

## Immunogen

A recombinant fragment (amino acids 73-187) from the human protein was used as the immunogen for the Granzyme B antibody.

## Storage

Store the Granzyme B antibody at 2-8°C (with azide) or aliquot and store at -20°C or colder (without azide).

## Alternate Names

Cathepsin G-like 1