



IL-12 Antibody (F51357)

Catalog No.	Formulation	Size
F51357-0.4ML	In 1X PBS, pH 7.4, with 0.09% sodium azide	0.4 ml
F51357-0.08ML	In 1X PBS, pH 7.4, with 0.09% sodium azide	0.08 ml


Bulk quote request

Availability	1-3 business days
Species Reactivity	Human
Format	Antigen affinity purified
Clonality	Polyclonal (rabbit origin)
Isotype	Rabbit Ig
Purity	Antigen affinity
UniProt	P29459
Localization	Cytoplasmic, secreted
Applications	Western Blot : 1:1000 IHC (Paraffin) : 1:10-1:50 Flow Cytometry : 1:10-1:50
Limitations	This IL-12 antibody is available for research use only.





IHC analysis of FFPE human breast carcinoma stained with IL-12 antibody



IL-12 antibody flow cytometry analysis of MDA-MB231 cells (green) compared to a <u>negative control</u> (blue). FITC-conjugated goat-anti-rabbit secondary Ab was used for the analysis.

Description

IL12A is a subunit of a cytokine that acts on T and natural killer cells, and has a broad array of biological activities. The cytokine is a disulfide-linked heterodimer composed of the 35-kD subunit encoded by this gene, and a 40-kD subunit that is a member of the cytokine receptor family. This cytokine is required for the T-cell-independent induction of interferon (IFN)-gamma, and is important for the differentiation of both Th1 and Th2 cells. The responses of lymphocytes to this cytokine are mediated by the activator of transcription protein STAT4. Nitric oxide synthase 2A (NOS2A/NOS2) is found to be required for the signaling process of this cytokine in innate immunity.

Application Notes

Titration of the IL-12 antibody may be required due to differences in protocols and secondary/substrate sensitivity.

Immunogen

A portion of amino acids 168-195 from human IL12A was used as the immunogen for this IL-12 antibody.

Storage

Aliquot the IL-12 antibody and store frozen at -20oC or colder. Avoid repeated freeze-thaw cycles.