

Insulin Receptor Antibody (F50643)


| Catalog No. | Formulation | Size |
|---------------|--|---------|
| F50643-0.2ML | In 1X PBS, pH 7.4, with 0.09% sodium azide | 0.2 ml |
| F50643-0.05ML | In 1X PBS, pH 7.4, with 0.09% sodium azide | 0.05 ml |

Bulk quote request

| | |
|-----------------------------|--|
| Availability | 1-3 business days |
| Species Reactivity | Human |
| Predicted Reactivity | Mouse, Rat |
| Format | Purified |
| Clonality | Polyclonal (rabbit origin) |
| Isotype | Rabbit Ig |
| Purity | Purified |
| UniProt | P06213 |
| Applications | Western Blot : 1:1000 IHC (Paraffin) : 1:50-1:100 |
| Limitations | This Insulin Receptor antibody is available for research use only. |

250
150
100
75
50
37
25
15

Western blot analysis of Insulin receptor antibody and SKBR-3 cell lysate. Expected size: ~156 kDa (precursor), ~95 kDa (b-subunit)



IHC analysis of FFPE human breast carcinoma tissue stained with the Insulin Receptor antibody

Description

INSR binds insulin and has a tyrosine-protein kinase activity. Autophosphorylation activates the kinase activity. This Type I membrane protein is composed of a tetramer of 2 alpha and 2 beta chains linked by disulfide bonds. The alpha chains contribute to the formation of the ligand-binding domain, while the beta chains carry the kinase domain. After being transported from the endoplasmic reticulum to the Golgi apparatus, the single glycosylated precursor is further glycosylated and then cleaved, followed by its transport to the plasma membrane.

Application Notes

Titration of the Insulin Receptor antibody may be required due to differences in protocols and secondary/substrate sensitivity.

Immunogen

A portion of amino acids 21-52 from the human protein was used as the immunogen for this Insulin Receptor antibody.

Storage

Aliquot the Insulin Receptor antibody and store frozen at -20°C or colder. Avoid repeated freeze-thaw cycles.