

Involucrin Antibody [clone SPM259] (V9036)


Catalog No.	Formulation	Size
V9036-100UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide	100 ug
V9036-20UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide	20 ug
V9036SAF-100UG	1 mg/ml in 1X PBS; BSA free, sodium azide free	100 ug
V9036IHC-7ML	Prediluted in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide; *For IHC use only*	7 ml



Citations (2)

[Bulk quote request](#)

Availability	1-3 business days
Species Reactivity	Human
Format	Purified
Host	Mouse
Clonality	Monoclonal (mouse origin)
Isotype	Mouse IgG1, kappa
Clone Name	SPM259
Purity	Protein G affinity chromatography
UniProt	P07476
Localization	Cytoplasmic
Applications	Immunohistochemistry (FFPE) : 0.1-0.2ug/ml for 30 min at RT
Limitations	This Involucrin antibody is available for research use only.



IHC: Formalin-fixed, paraffin-embedded human skin stained with Involucrin antibody (SPM259)

Description

Involucrin is expressed in a range of stratified squamous epithelia, including the cornea, which lacks a distinct cornified layer. In normal epidermis, it is first expressed in the upper spinous layers, and in keratinocyte cultures, all cells that have left the basal layer express it. Involucrin expression is altered in pathological conditions: in psoriasis and other benign epidermal hyperplasias, involucrin expression begins closer to the basal layer than normal; expression is abnormal in squamous cell carcinomas and premalignant lesions, and is reduced in severe dysplasias of the larynx and cervix.

Application Notes

The optimal dilution of the Involucrin antibody for each application should be determined by the researcher.

1. Staining of formalin-fixed tissues is enhanced by digestion with Trypsin or Protease XXV at 1mg/ml PBS for 5 min at 37oC. Enzyme digestion is better than citrate buffer based epitope unmasking.
2. The prediluted format is supplied in a dropper bottle and is optimized for use in IHC. After epitope retrieval step (if required), drip mAb solution onto the tissue section and incubate at RT for 30 min.

Immunogen

Purified involucrin from human keratinocytes was used as the immunogen for this Involucrin antibody. Its epitope maps between amino acids 421-568 of human involucrin.

Storage

Store the Involucrin antibody at 2-8oC (with azide) or aliquot and store at -20oC or colder (without azide).