


IQCJ Antibody / IQ domain-containing protein J (F54564)


| Catalog No. | Formulation | Size |
|---------------|--------------------------------------------|---------|
| F54564-0.4ML | In 1X PBS, pH 7.4, with 0.09% sodium azide | 0.4 ml |
| F54564-0.08ML | In 1X PBS, pH 7.4, with 0.09% sodium azide | 0.08 ml |

[Bulk quote request](#)


| | |
|---------------------------|-----------------------------------------------------------------------------------------------------------|
| Availability | 1-3 business days |
| Species Reactivity | Human, Mouse |
| Format | Purified |
| Host | Rabbit |
| Clonality | Polyclonal (rabbit origin) |
| Isotype | Rabbit Ig |
| Purity | Antigen affinity purified |
| UniProt | Q1A5X6 |
| Applications | Flow Cytometry : 1:25 (1x10e6 cells) Immunohistochemistry (FFPE) : 1:25 Western Blot : 1:500-1:2000 |
| Limitations | This IQCJ antibody is available for research use only. |




Western blot testing of mouse NIH 3T3 cell lysate with IQCJ antibody. Predicted molecular weight ~18 kDa.



Western blot testing of human K562 cell lysate with IQCJ antibody. Predicted molecular weight ~18 kDa.



IHC testing of FFPE human brain tissue with IQCJ antibody. HIER: steam section in pH6 citrate buffer for 20 min and allow to cool prior to staining.



Flow cytometry testing of human K562 cells with IQCJ antibody; Blue=isotype control, Green= IQCJ antibody.

Description

Alternate names: IQ domain-containing protein J

Application Notes

The stated application concentrations are suggested starting points. Titration of the IQCJ antibody may be required due to differences in protocols and secondary/substrate sensitivity.

Immunogen

A portion of amino acids 98-126 from the human protein was used as the immunogen for the IQCJ antibody.

Storage

Aliquot the IQCJ antibody and store frozen at -20oC or colder. Avoid repeated freeze-thaw cycles.