


## NADH-ubiquinone oxidoreductase chain 5 Antibody / MT-ND5 (F54903)


Catalog No.	Formulation	Size
F54903-0.4ML	In 1X PBS, pH 7.4, with 0.09% sodium azide	0.4 ml
F54903-0.08ML	In 1X PBS, pH 7.4, with 0.09% sodium azide	0.08 ml

**Bulk quote request**


<b>Availability</b>	1-3 business days
<b>Species Reactivity</b>	Human
<b>Format</b>	Purified
<b>Host</b>	Rabbit
<b>Clonality</b>	Polyclonal (rabbit origin)
<b>Isotype</b>	Rabbit Ig
<b>Purity</b>	Antigen affinity purified
<b>UniProt</b>	P03915
<b>Localization</b>	Cytoplasmic
<b>Applications</b>	Western Blot : 1:500-1:1000 Flow Cytometry : 1:10-1:50 (1x10 <sup>6</sup> cells) Immunohistochemistry (FFPE) : 1:100
<b>Limitations</b>	This NADH-ubiquinone oxidoreductase chain 5 antibody is available for research use only.




IHC testing of FFPE human liver tissue with NADH-ubiquinone oxidoreductase chain 5 antibody. HIER: steam section in pH6 citrate buffer for 20 min and allow to cool prior to staining.



IHC testing of FFPE human brain tissue with NADH-ubiquinone oxidoreductase chain 5 antibody. HIER: steam section in pH6 citrate buffer for 20 min and allow to cool prior to staining.



Western blot testing of human 1) CCRF-CEM and 2) Jurkat cell lysate with NADH-ubiquinone oxidoreductase chain 5 antibody. Expected molecular weight: 65-70 kDa.



Flow cytometry testing of human CCRF-CEM cells with NADH-ubiquinone oxidoreductase chain 5 antibody; Blue=isotype control, Green= NADH-ubiquinone oxidoreductase chain 5 antibody.

## Description

Core subunit of the mitochondrial membrane respiratory chain NADH dehydrogenase (Complex I) that is believed to belong to the minimal assembly required for catalysis. Complex I functions in the transfer of electrons from NADH to the respiratory chain. The immediate electron acceptor for the enzyme is believed to be ubiquinone.

## Application Notes

The stated application concentrations are suggested starting points. Titration of the NADH-ubiquinone oxidoreductase chain 5 antibody may be required due to differences in protocols and secondary/substrate sensitivity.

## Immunogen

A portion of amino acids 544-570 from the human protein was used as the immunogen for the NADH-ubiquinone oxidoreductase chain 5 antibody.

## Storage

Aliquot the NADH-ubiquinone oxidoreductase chain 5 antibody and store frozen at -20°C or colder. Avoid repeated freeze-thaw cycles.

