


NANOG Antibody (F53329)


Catalog No.	Formulation	Size
F53329-0.4ML	In 1X PBS, pH 7.4, with 0.09% sodium azide	0.4 ml
F53329-0.08ML	In 1X PBS, pH 7.4, with 0.09% sodium azide	0.08 ml

[Bulk quote request](#)

Availability	1-3 business days
Species Reactivity	Human
Format	Antigen affinity purified
Host	Rabbit
Clonality	Polyclonal (rabbit origin)
Isotype	Rabbit Ig
Purity	Antigen affinity
UniProt	Q9H9S0
Applications	Western Blot : 1:1000-2000
Limitations	This NANOG antibody is available for research use only.



Western blot testing of NANOG antibody at 1:2000 dilution + human brain lysate;
Predicted molecular weight: 35-45 kDa.



Description

Transcription regulator involved in inner cell mass and embryonic stem (ES) cells proliferation and self-renewal. Imposes pluripotency on ES cells and prevents their differentiation towards extraembryonic endoderm and trophectoderm lineages. Blocks bone morphogenetic protein-induced mesoderm differentiation of ES cells by physically interacting with SMAD1 and interfering with the recruitment of coactivators to the active SMAD transcriptional complexes. Acts as a transcriptional activator or repressor. Binds optimally to the DNA consensus sequence 5'-TAAT[GT][GT]-3' or 5'-[CG][GA][CG]C[GC]ATTAN[GC]-3'. Able to autorepress its expression in differentiating (ES) cells: binds to its own promoter following interaction with ZNF281/ZFP281, leading to recruitment of the NuRD complex and subsequent repression of expression. When overexpressed, promotes cells to enter into S phase and proliferation.

Application Notes

Titration of the NANOG antibody may be required due to differences in protocols and secondary/substrate sensitivity.

Immunogen

This NANOG antibody was produced from a rabbit immunized with a KLH conjugated synthetic peptide between 155-189 amino acids from the Central region of human NANOG.

Storage

Aliquot the NANOG antibody and store frozen at -20oC or colder. Avoid repeated freeze-thaw cycles.