


NuMA Antibody [clone SPM300] (V2768)


| Catalog No. | Formulation | Size |
|----------------|--|--------|
| V2768-100UG | 0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide | 100 ug |
| V2768-20UG | 0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide | 20 ug |
| V2768SAF-100UG | 1 mg/ml in 1X PBS; BSA free, sodium azide free | 100 ug |

Bulk quote request

| | |
|---------------------------|---|
| Availability | 1-3 business days |
| Species Reactivity | Human |
| Format | Purified |
| Clonality | Monoclonal (mouse origin) |
| Isotype | Mouse IgM, kappa |
| Clone Name | SPM300 |
| Purity | PEG precipitation |
| UniProt | Q14980 |
| Localization | Nuclear |
| Applications | Immunohistochemistry (FFPE) : 1-2ug/ml for 30 min at RT |
| Limitations | This NuMA antibody is available for research use only. |



IHC analysis of formalin-fixed, paraffin-embedded human tonsil stained with NuMA antibody (clone SPM300).



SDS-PAGE analysis of purified, BSA-free NuMA antibody (clone SPM300) as confirmation of integrity and purity.

Description

Recognizes a phosphorylated protein of 228kDa, identified as nuclear mitotic apparatus protein (NuMA). NuMA is intranuclear protein and present in nucleus during interphase. At the onset of mitosis, it redistributes from the nucleus to two centrosomal structures that later will become part of the mitotic spindle pole. After anaphase, the protein redistributes from the spindle polar region into reforming nucleus. NuMA is an essential protein during mitosis for the terminal phases of chromosome separation and/or nuclear reassembly. Recently a study shows that NuMA is cleaved to a 180 to 200kDa during apoptosis. Chromosomal translocation of this gene with the RARA (retinoic acid receptor, alpha) gene on chromosome 17 has been detected in patients with acute promyelocytic leukemia.

Application Notes

Optimal dilution of the NuMA antibody should be determined by the researcher.

1. Staining of formalin-fixed tissues requires boiling tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 10-20 min followed by cooling at RT for 20 min.

Immunogen

Colon carcinoma 174T cells were used as the immunogen for the NuMA antibody.

Storage

Store the NuMA antibody at 2-8oC (with azide) or aliquot and store at -20oC or colder (without azide).