


Otoferlin Antibody / OTOF (R32323)


Catalog No.	Formulation	Size
R32323	0.5mg/ml if reconstituted with 0.2ml sterile DI water	100 ug

[Bulk quote request](#)


Availability	1-3 business days
Species Reactivity	Mouse, Rat
Format	Antigen affinity purified
Host	Rabbit
Clonality	Polyclonal (rabbit origin)
Isotype	Rabbit IgG
Purity	Antigen affinity
Buffer	Lyophilized from 1X PBS with 2% Trehalose
UniProt	Q9HC10
Localization	Cytoplasmic, cell membrane
Applications	Western Blot : 0.5-1ug/ml Immunohistochemistry (FFPE) : 2-5ug/ml
Limitations	This Otoferlin antibody is available for research use only.



IHC testing of FFPE mouse brain tissue with Otoferlin antibody. HIER: Boil the paraffin sections in pH 8 EDTA buffer for 20 minutes and allow to cool prior to staining.



IHC testing of FFPE rat brain tissue with Otoferlin antibody. HIER: Boil the paraffin sections in pH 8 EDTA buffer for 20 minutes and allow to cool prior to staining.



Western blot testing of 1) rat brain and 2) mouse brain tissue lysate with Otofelin antibody. Predicted molecular weight: 226-227 kDa (long form, two isoforms), 140-149 kDa (short form, three isoforms).

Description

Otofelin is a protein that in humans is encoded by the OTOF gene. Mutations in this gene are a cause of neurosensory nonsyndromic recessive deafness, DFNB9. The short form of the encoded protein has three C2 domains, a single carboxy-terminal transmembrane domain found also in the *C. elegans* spermatogenesis factor FER-1 and human dysferlin, while the long form has six C2 domains. The homology suggests that this protein may be involved in vesicle membrane fusion. Several transcript variants encoding multiple isoforms have been found for this gene.

Application Notes

Optimal dilution of the Otofelin antibody should be determined by the researcher.

Immunogen

Amino acids QIWDADHFSADDFLGAIELDLNRFPRGAKTAKQ of human OTOF were used as the immunogen for the Otofelin antibody.

Storage

After reconstitution, the Otofelin antibody can be stored for up to one month at 4°C. For long-term, aliquot and store at -20°C. Avoid repeated freezing and thawing.