



p62 Antibody (F48010)

Catalog No.	Formulation	Size
F48010-0.4ML	In 1X PBS, pH 7.4, with 0.09% sodium azide	0.4 ml
F48010-0.08ML	In 1X PBS, pH 7.4, with 0.09% sodium azide	0.08 ml


Bulk quote request


Availability	1-3 business days
Species Reactivity	Human, African Green Monkey
Predicted Reactivity	Mouse, Rat
Format	Antigen affinity purified
Clonality	Polyclonal (rabbit origin)
Isotype	Rabbit Ig
Purity	Antigen affinity
UniProt	Q13501
Applications	Western Blot : 1:1000 IHC (Paraffin) : 1:50-1:100 Immunofluorescence : 1:50-1:100
Limitations	This p62 antibody is available for research use only.







Fluorescent image of U251 cells stained with p62 antibody at 1:200. Immunoreactivity is localized to autophagic vacuoles in the cytoplasm.







IHC analysis of FFPE human hepatocarcinoma tissue stained with the p62 antibody



Immunofluorescence staining of p62 antibody on Methanol-fixed and PFA fixed HeLa cells. Data courtesy of Dr. Eeva-Liisa Eskelinen, University of Helsinki, Finland.

Description

SQSTM1/p62 is an adapter protein which binds ubiquitin and may regulate the activation of NFKB1 by TNF-alpha, nerve growth factor (NGF) and interleukin-1. This protein may play a role in titin/TTN downstream signaling in muscle cells, and may also regulate signaling cascades through ubiquitination. This protein is involved in cell differentiation, apoptosis, immune response and regulation of K(+) channels. SQSTM1/p62 also appears to play a role in macroautophagic removal of intracellular protein aggregates. Cellular depletion studies of SQSTM1/p62 have indicated a role for association with LC3 and aggregate proteins in order to facilitate normal formation of the autophagosome.

Application Notes

Titration of the p62 antibody may be required due to differences in protocols and secondary/substrate sensitivity.

Immunogen

A portion of amino acids 317-346 from the human protein was used as the immunogen for this p62 antibody.

Storage

Aliquot the p62 antibody and store frozen at -20oC or colder. Avoid repeated freeze-thaw cycles.		