


## p70S6K Antibody (F48468)


Catalog No.	Formulation	Size
F48468-0.2ML	In 1X PBS, pH 7.4, with 0.09% sodium azide	0.2 ml
F48468-0.05ML	In 1X PBS, pH 7.4, with 0.09% sodium azide	0.05 ml

**Bulk quote request**


<b>Availability</b>	1-3 business days
<b>Species Reactivity</b>	Human
<b>Predicted Reactivity</b>	Bovine, Mouse, Rabbit, Rat
<b>Format</b>	Antigen affinity purified
<b>Clonality</b>	Polyclonal (rabbit origin)
<b>Isotype</b>	Rabbit Ig
<b>Purity</b>	Antigen affinity
<b>UniProt</b>	P23443
<b>Applications</b>	Western Blot : 1:1000 IHC (Paraffin) : 1:10-1:50
<b>Limitations</b>	This p70S6K antibody is available for research use only.



Western blot analysis of p70S6K antibody and TNF alpha activated HeLa lysate; both phospho and non-phos S6K can be detected with this Ab. Predicted molecular weight: 60-70 kDa



Western blot analysis of p70S6K antibody and 293 cell lysate (2 ug/lane) either nontransfected (Lane 1) or transiently transfected with the RPS6KB1 gene (2).



IHC analysis of FFPE human breast carcinoma tissue stained with p70S6K antibody

## Description

RPS6kB1 is a serine/threonine-protein kinase that acts downstream of mTOR signaling in response to growth factors and nutrients to promote cell proliferation, cell growth and cell cycle progression. Regulates protein synthesis through phosphorylation of EIF4B, RPS6 and EEF2K, and contributes to cell survival by repressing the pro-apoptotic function of BAD. Under conditions of nutrient depletion, the inactive form associates with the EIF3 translation initiation complex. Upon mitogenic stimulation, phosphorylation by the mammalian target of rapamycin complex 1 (mTORC1) leads to dissociation from the EIF3 complex and activation. The active form then phosphorylates and activates several substrates in the pre-initiation complex, including the EIF2B complex and the cap-binding complex component EIF4B. [UniProt]

## Application Notes

The stated application concentrations are suggested starting amounts. Titration of the p70S6K antibody may be required due to differences in protocols and secondary/substrate sensitivity.

## Immunogen

A portion of amino acids 425-454 from human RPS6KB1 was used as the immunogen for this p70S6K antibody.

## Storage

Aliquot the p70S6K antibody and store frozen at -20°C or colder. Avoid repeated freeze-thaw cycles.