

Perlecan Antibody (HSPG) [clone A7L6] (V2600)


Catalog No.	Formulation	Size
V2600-100UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide	100 ug
V2600-20UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide	20 ug
V2600SAF-100UG	1 mg/ml in 1X PBS; BSA free, sodium azide free	100 ug
V2600IHC-7ML	Prediluted in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide; *For IHC use only*	7 ml




Citations (10)

[Bulk quote request](#)

Availability	1-3 business days
Species Reactivity	Human, Mouse
Format	Purified
Clonality	Monoclonal (rat origin)
Isotype	Rat IgG2a, kappa
Clone Name	A7L6
Purity	Protein G affinity chromatography
UniProt	P98160
Localization	Basement membrane
Applications	Flow Cytometry : 0.5-1ug/10 ⁶ cells Immunofluorescence : 0.5-1ug/ml Immunohistochemistry (FFPE) : 1-2ug/ml for 30 min at RT
Limitations	This Perlecan antibody is available for research use only.



IHC: Formalin-fixed, paraffin-embedded human testicular carcinoma stained with Perlecan antibody (A7L6).



SDS-PAGE Analysis of Purified, BSA-Free Perlecan Antibody (clone A7L6).
Confirmation of Integrity and Purity of the Antibody.

Description

This mAb specifically precipitates heterogeneous material of high MW, identified as perlecan, a major heparan-sulfate proteoglycan (HSPG) within all basement membranes and cell surfaces. It does not cross-react with laminin, fibronectin, or dermatan sulfate proteoglycan. Because of perlecan's strategic location and ability to store and protect growth factors, it has been strongly implicated in the control of tumor cell growth and metastatic behavior. Perlecan possesses angiogenic and growth-promoting attributes primarily by acting as a co-receptor for basic fibroblast growth factor (FGF-2). Suppression of perlecan causes substantial inhibition of neoplastic growth and neovascularization. Thus, perlecan is a potent inducer of neoplasm growth and angiogenesis in vivo and therapeutic interventions targeting this key modulator of tumor progression may improve neoplastic treatment.

Application Notes

Optimal dilution of the Perlecan antibody should be determined by the researcher.

1. Staining of formalin-fixed tissues requires boiling tissue sections in 10mM Citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 min
2. The prediluted format is supplied in a dropper bottle and is optimized for use in IHC. After epitope retrieval step (if required), drip mAb solution onto the tissue section and incubate at RT for 30 min.

Immunogen

A murine EHS laminin preparation was used as the immunogen for the Perlecan antibody.

Storage

Store the Perlecan antibody at 2-8°C (with azide) or aliquot and store at -20°C or colder (without azide).