


PIASg Antibody / PIAS4 (R31339)

Catalog No.	Formulation	Size
R31339	0.5mg/ml if reconstituted with 0.2ml sterile DI water	100 ug

Bulk quote request

Availability	1-3 business days
Species Reactivity	Human, Mouse, Rat
Format	Antigen affinity purified
Host	Rabbit
Clonality	Polyclonal (rabbit origin)
Isotype	Rabbit IgG
Purity	Antigen affinity
Buffer	Lyophilized from 1X PBS with 2.5% BSA and 0.025% sodium azide/thimerosal
UniProt	Q8N2W9
Applications	Western Blot : 0.5-1ug/ml
Limitations	This PIASg antibody is available for research use only.



Western blot testing of PIASg antibody and Lane 1: human placenta; 2: A431 lysate.
Expected/observed size ~56 kDa.

Description

Protein inhibitor of activated STAT protein 4 (PIAS4) or Protein inhibitor of activated STAT protein gamma (PIASg or PIASy) is an enzyme that in humans is encoded by the PIAS4 gene. This gene is mapped to 19p13.3. This gene plays a crucial role as a transcriptional coregulation in various cellular pathways, including the STAT pathway, the p53/TP53 pathway, the Wnt pathway and the steroid hormone signaling pathway. PIASg functions as an E3-type small ubiquitin-like modifier (SUMO) ligase, stabilizing the interaction between UBE2I and the substrate, and as a SUMO-tethering factor. This gene involved in gene silencing.

Application Notes

The stated application concentrations are suggested starting amounts. Titration of the PIASg antibody may be required due to differences in protocols and secondary/substrate sensitivity.

Immunogen

An amino acid sequence from the C-terminus of human E3 SUMO-protein ligase PIASg (HPELCKALVKEKLRLD) was used as the immunogen for this PIASg antibody (100% homologous in human, mouse and rat).

Storage

After reconstitution, the PIASg antibody can be stored for up to one month at 4°C. For long-term, aliquot and store at -20°C. Avoid repeated freezing and thawing.