


## Pro-MCH Antibody / PMCH (F54416)


Catalog No.	Formulation	Size
F54416-0.4ML	In 1X PBS, pH 7.4, with 0.09% sodium azide	0.4 ml
F54416-0.08ML	In 1X PBS, pH 7.4, with 0.09% sodium azide	0.08 ml

[Bulk quote request](#)


<b>Availability</b>	1-3 business days
<b>Species Reactivity</b>	Human, Mouse
<b>Format</b>	Purified
<b>Clonality</b>	Polyclonal (rabbit origin)
<b>Isotype</b>	Rabbit Ig
<b>Purity</b>	Antigen affinity purified
<b>UniProt</b>	P20382
<b>Applications</b>	Western Blot : 1:500-1:2000 Flow Cytometry : 1:25 (1x10e6 cells) Immunohistochemistry (FFPE) : 1:25
<b>Limitations</b>	This Pro-MCH antibody is available for research use only.




Western blot testing of mouse brain lysate with Pro-MCH antibody.




Western blot testing of human K562 cell lysate with Pro-MCH antibody.



Western blot testing of 1) non-transfected and 2) transfected 293 cell lysate with Pro-MCH antibody.



IHC testing of FFPE human testis tissue with Pro-MCH antibody. HIER: steam section in pH6 citrate buffer for 20 min and allow to cool prior to staining.



Flow cytometry testing of human K562 cells with Pro-MCH antibody; Blue=isotype control, Green= Pro-MCH antibody.

## Description

The melanin-concentrating hormone (MCH) is a cyclic neuropeptide isolated initially from salmon pituitary gland and later from rat hypothalamus (summarized by Nahon et al., 1992 [PubMed 1572663]). In mammals, MCH perikarya are confined largely to the lateral hypothalamus and zona incerta area with extensive neuronal projections throughout the brain, including the neurohypophysis. The anatomic distribution suggests a neurotransmitter or neuromodulator role for MCH in a broad array of neuronal functions directed toward the regulation of goal-directed behavior, such as food intake, and general arousal. MCH and 2 other putative neuropeptides, NEI and NGE, are encoded by the same precursor and appear colocalized in nerve cells and in many instances within the projections. The precursor is designated pro-melanin-concentrating hormone (PMCH).

## Application Notes

The stated application concentrations are suggested starting points. Titration of the Pro-MCH antibody may be required due to differences in protocols and secondary/substrate sensitivity.

## Immunogen

A portion of amino acids 94-122 from the human protein was used as the immunogen for the Pro-MCH antibody.

## Storage

Aliquot the Pro-MCH antibody and store frozen at -20oC or colder. Avoid repeated freeze-thaw cycles.

