

S100A9 Antibody [clone 47-8D3] (V2852)


Catalog No.	Formulation	Size
V2852-100UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide	100 ug
V2852-20UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide	20 ug
V2852SAF-100UG	1 mg/ml in 1X PBS; BSA free, sodium azide free	100 ug




Citations (4)

[Bulk quote request](#)

Availability	1-3 business days
Species Reactivity	Human, Rat
Format	Purified
Clonality	Monoclonal (mouse origin)
Isotype	Mouse IgG1, kappa
Clone Name	47-8D3
Purity	Protein G affinity chromatography
UniProt	P06702
Localization	Cytoplasmic
Applications	Flow Cytometry : 0.5-1ug/10 ⁶ cells Immunofluorescence : 0.5-1ug/ml Immunohistochemistry (FFPE) : 0.5-1ug/ml for 30 min at RT
Limitations	This S100A9 antibody is available for research use only.



Western blot testing of human PBM lysate with S100A9 antibody (clone 47-8D3).
Expected/observed molecular weight ~14kDa.



SDS-PAGE Analysis of Purified, BSA-Free S100A9 Antibody (clone 47-8D3).
Confirmation of Integrity and Purity of the Antibody.

Description

This mAb stains the cytoplasm of macrophages and histiocytes in hematopoietic organs, Kupffer's cells of the liver and Langerhan's cells of the skin. It also stains the mantle zone B-lymphocytes of the lymph node and spleen, spermatogonia, and chief cells of the stomach. S100A9 is expressed by macrophages in acutely inflamed tissues and in chronic inflammation. It is detected in peripheral blood leukocytes, in neutrophils and granulocytes. It is present at sites of vascular inflammation. S100A9 is also expressed in epithelial cells constitutively or induced during dermatoses. S100A9 is a Calcium-binding protein. It has antimicrobial activity towards bacteria and fungi. It is important for resistance to invasion by pathogenic bacteria. It up-regulates transcription of genes that are under the control of NF-kappa-B. S100A9 plays a role in the development of endotoxic shock in response to bacterial lipopolysaccharide (LPS). It promotes tubulin polymerization when unphosphorylated. It also promotes phagocyte migration and infiltration of granulocytes at sites of wounding. It plays a role as a pro-inflammatory mediator in acute and chronic inflammation and up-regulates the release of IL8 and cell-surface expression of ICAM1.

Application Notes

Optimal dilution of the S100A9 antibody should be determined by the researcher.

1. For staining of formalin-fixed, paraffin-embedded tissues, digest with Trypsin at 1mg/ml PBS, 15 min at RT

Immunogen

Human peripheral blood monocyte components were used as the immunogen for the S100A9 antibody.

Storage

Store the S100A9 antibody at 2-8°C (with azide) or aliquot and store at -20°C or colder (without azide).

Alternate Names

Calprotectin L1H subunit , MRP14, Calgranulin B