


SPHK1 Antibody (F50216)


Catalog No.	Formulation	Size
F50216-0.4ML	In 1X PBS, pH 7.4, with 0.09% sodium azide	0.4 ml
F50216-0.08ML	In 1X PBS, pH 7.4, with 0.09% sodium azide	0.08 ml

Bulk quote request

Availability	1-3 business days
Species Reactivity	Human
Format	Purified
Host	Rabbit
Clonality	Polyclonal (rabbit origin)
Isotype	Rabbit Ig
Purity	Purified
UniProt	Q9NYA1
Applications	Western Blot : 1:500
Limitations	This SPHK1 antibody is available for research use only.



The SpHK1 antibody used in western blot (lane 2) to detect c-myc-tagged SpHK1 in transfected 293 cell lysate (a c-myc antibody is used as control in Lane 1). Predicted molecular weight: ~43/51/44kDa (isoforms 1/2/3).



Western blot analysis of SPHK1 antibody and 293T cell lysate either nontransfected (Lane 1) or transiently transfected (2) with the SPHK1 gene. Predicted molecular weight: ~43/51/44kDa (isoforms 1/2/3).



Western blot analysis of lysate from A431 cell line using SPHK1 antibody at 1:1000.
Predicted molecular weight: ~43/51/44kDa (isoforms 1/2/3).

Description

Sphingosine Kinase (SphK) catalyzes the phosphorylation of the lipid sphingosine, creating the bioactive lipid sphingosine-1-phosphate (S1P). S1P subsequently signals through cell surface G protein-coupled receptors, as well as intracellularly, to modulate cell proliferation, survival, motility and differentiation. SphK is an important signaling enzyme which is activated by diverse agents, including growth factors that signal through receptor tyrosine kinases, agents activating G protein-coupled receptors, and immunoglobulin receptors. Two SphK isotypes, SphK-1 and SphK-2, have been cloned, and both isotypes are ubiquitously expressed. SphK-1 has been shown to mediate cell growth, prevention of apoptosis, and cellular transformation, and is upregulated in a variety of human tumors. In contrast, SphK-2 increases apoptosis, and may be responsible for phosphorylating and activating the immunosuppressive drug FTY720.

Application Notes

Titration of the SPHK1 antibody may be required due to differences in protocols and secondary/substrate sensitivity.

Immunogen

A portion of amino acids 1-30 from the human protein was used as the immunogen for this SPHK1 antibody.

Storage

Aliquot the SPHK1 antibody and store frozen at -20oC or colder. Avoid repeated freeze-thaw cycles.