


TRIM65 Antibody (F54616)


| Catalog No. | Formulation | Size |
|---------------|--|---------|
| F54616-0.4ML | In 1X PBS, pH 7.4, with 0.09% sodium azide | 0.4 ml |
| F54616-0.08ML | In 1X PBS, pH 7.4, with 0.09% sodium azide | 0.08 ml |

Bulk quote request

| | |
|---------------------------|--|
| Availability | 1-3 business days |
| Species Reactivity | Human |
| Format | Purified |
| Host | Rabbit |
| Clonality | Polyclonal (rabbit origin) |
| Isotype | Rabbit Ig |
| Purity | Antigen affinity purified |
| UniProt | Q6PJ69 |
| Localization | Cytoplasmic, nuclear |
| Applications | Flow Cytometry : 1:25 (1x10e6 cells) Immunofluorescence : 1:25 Immunohistochemistry (FFPE) : 1:25 Western Blot : 1:500-1:2000 |
| Limitations | This TRIM65 antibody is available for research use only. |




IHC testing of FFPE human prostate carcinoma tissue with TRIM65 antibody. HIER: steam section in pH6 citrate buffer for 20 min and allow to cool prior to staining.



Immunofluorescent staining of human NCI-H460 cells with TRIM65 antibody (green), DAPI nuclear stain (blue) and anti-Actin (red).

kDa
130
72
55 ->
36
28
17

Western blot testing of human NCI-H460 cell lysate with TRIM65 antibody. Predicted molecular weight ~57 kDa.



Flow cytometry testing of human NCI-H460 cells with TRIM65 antibody; Blue=isotype control, Green= TRIM65 antibody.

Description

TRIM65 belongs to the TRIM/RBCC family. It contains one B box-type zinc finger, one B30.2/SPRY domain and one RING-type zinc finger.

Application Notes

The stated application concentrations are suggested starting points. Titration of the TRIM65 antibody may be required due to differences in protocols and secondary/substrate sensitivity.

Immunogen

A portion of amino acids 327-355 from the human protein was used as the immunogen for the TRIM65 antibody.

Storage

Aliquot the TRIM65 antibody and store frozen at -20oC or colder. Avoid repeated freeze-thaw cycles.