


von Willebrand Factor Antibody [clone WFA52-2] (V7082)


Catalog No.	Formulation	Size
V7082-100UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide	100 ug
V7082-20UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide	20 ug
V7082SAF-100UG	1 mg/ml in 1X PBS; BSA free, sodium azide free	100 ug

[Bulk quote request](#)

Species Reactivity	Human
Format	Purified
Host	Mouse
Clonality	Monoclonal (mouse origin)
Isotype	Mouse IgG1, kappa
Clone Name	WFA52-2
Purity	Protein G purified
Gene ID	7450
Localization	Cytoplasmic
Applications	Immunohistochemistry (FFPE) : 1-2ug/ml for 30 min at RT
Limitations	This von Willebrand Factor antibody is available for research use only.



IHC testing of FFPE human pancreas and von Willebrand Factor antibody (clone WFA52-2). Required HIER: boil tissue sections in pH 9 10mM Tris with 1mM EDTA for 10-20 min.



IHC staining of human bone marrow with von Willebrand Factor antibody (clone WFA52-2). Required HIER: boil tissue sections in pH 9 10mM Tris with 1mM EDTA for 10-20 min.

Description

von Willebrand Factor (vWF) is a glycoprotein produced by bone marrow cells and endothelial cells lining the inside surface of blood vessels. Its primary function is platelet adhesion, binding to Factor VIII, collagen and platelets, to coagulate blood at the site of wounding. The enzyme VWFCP, or vWF-cleaving protease, facilitates clotting by cutting the protein into subunits, increasing its binding capacity.

Deficiency or dysfunction of the protein increases the tendency of wounds to bleed, or to bleed more. $\sim 1/2$ Over 300 gene mutations have been identified and classified into three types. $\sim 1/2$ Type 1 von Willebrand Factor disease is characterized by reduced amounts in the bloodstream, Type 2 by reduced binding ability and Type 3 by a nonfunctional protein.

Application Notes

Variations in protocols, secondaries and substrates may require the von Willebrand Factor antibody to be titrated for optimal performance.

Immunogen

A recombinant human protein fragment was used as the immunogen for this von Willebrand Factor antibody.

Storage

Store the von Willebrand Factor antibody at 2-8°C (with azide) or aliquot and store at -20°C or colder (without azide).