


## CKS2 Antibody / Cyclin-dependent kinases regulatory subunit 2 (N-Terminal Region) (F54161)

| Catalog No.   | Formulation                                | Size    |
|---------------|--------------------------------------------|---------|
| F54161-0.2ML  | In 1X PBS, pH 7.4, with 0.09% sodium azide | 0.2 ml  |
| F54161-0.05ML | In 1X PBS, pH 7.4, with 0.09% sodium azide | 0.05 ml |

[Bulk quote request](#)

|                      |                                                        |
|----------------------|--------------------------------------------------------|
| Availability         | 1-3 business days                                      |
| Species Reactivity   | Human                                                  |
| Predicted Reactivity | Bovine, Mouse                                          |
| Format               | Antigen affinity purified                              |
| Host                 | Rabbit                                                 |
| Clonality            | Polyclonal (rabbit origin)                             |
| Isotype              | Rabbit Ig                                              |
| Purity               | Antigen affinity                                       |
| UniProt              | P33552                                                 |
| Applications         | Western Blot : 1:1000-1:2000                           |
| Limitations          | This CKS2 antibody is available for research use only. |



Western blot testing of human 1) Caco-2, 2) RMPI-8226 and 3) SH-SY5Y cell lysate with CKS2 antibody at 1:2000. Predicted molecular weight ~10 kDa.

### Description

Cyclin-dependent kinases regulatory subunit 2 (CKS2) binds to the catalytic subunit of the cyclin dependent kinases and is essential for their biological function.

### Application Notes

The stated application concentrations are suggested starting points. Titration of the CKS2 antibody may be required due to differences in protocols and secondary/substrate sensitivity.

## **Immunogen**

A portion of amino acids 6-40 from human CKS2 was used as the immunogen for the CKS2 antibody.

## **Storage**

Aliquot the CKS2 antibody and store frozen at -20oC or colder. Avoid repeated freeze-thaw cycles.