


ADO Antibody / Aminoethanethiol dioxygenase (R32500)




Catalog No.	Formulation	Size
R32500	0.5mg/ml if reconstituted with 0.2ml sterile DI water	100 ug



[Bulk quote request](#)

Availability	1-3 business days
Species Reactivity	Human, Mouse, Rat
Format	Antigen affinity purified
Clonality	Polyclonal (rabbit origin)
Isotype	Rabbit IgG
Purity	Antigen affinity
Buffer	Lyophilized from 1X PBS with 2.5% BSA and 0.025% sodium azide
UniProt	Q96SZ5
Localization	Cytoplasmic, membranous
Applications	Western blot : 0.5-1ug/ml Immunofluorescence : 2-4ug/ml Flow cytometry : 1-3ug/million cells
Limitations	This ADO antibody is available for research use only.



Immunofluorescent staining of FFPE human A549 cells with ADO antibody (green) at 2ug/ml and DAPI nuclear stain (blue). HIER: steam section in pH6 citrate buffer for 20 min.





Description

Human thiol dioxygenases include cysteine dioxygenase (CDO) and cysteamine (2-aminoethanethiol) dioxygenase (ADO). CDO adds 2 oxygen atoms to free cysteine, whereas ADO adds 2 oxygen atoms to free cysteamine to form hypotaurine. It is demonstrated that mouse Ado has strong and specific dioxygenase activity in vitro towards cysteamine but not cysteine. Recombinant Ado was shown to bind iron. Overexpression of Ado in HepG2/C3A cells increased the production of hypotaurine from cysteamine. Similar results were found with human ADO. When endogenous expression of ADO was reduced by RNA-mediated interference, hypotaurine production decreased. It is also noted that the demonstration of high levels of ADO in brain challenges the previous assumption that most of the taurine in the brain is a consequence of CDO activity.

Application Notes

Differences in protocols and secondary/substrate sensitivity may require the ADO antibody to be titrated for optimal performance.

Immunogen

Amino acids E49-E261 from the human protein were used as the immunogen for the ADO antibody.

Storage

After reconstitution, the ADO antibody can be stored for up to one month at 4oC. For long-term, aliquot and store at -20oC. Avoid repeated freezing and thawing.