


## ACVRL1 Antibody / ALK1 (RQ6410)


Catalog No.	Formulation	Size
RQ6410	0.5mg/ml if reconstituted with 0.2ml sterile DI water	100 ug

**Bulk quote request**


<b>Availability</b>	1-3 business days
<b>Species Reactivity</b>	Human, Mouse, Rat
<b>Format</b>	Purified
<b>Clonality</b>	Polyclonal (rabbit origin)
<b>Isotype</b>	Rabbit IgG
<b>Purity</b>	Antigen affinity purified
<b>Buffer</b>	Lyophilized from 1X PBS with 2% Trehalose
<b>UniProt</b>	P37023
<b>Localization</b>	Cytoplasmic, cell membrane
<b>Applications</b>	Western blot : 0.5-1ug/ml Immunohistochemistry (FFPE) : 2-5ug/ml Flow cytometry : 1-3ug/million cells Direct ELISA : 0.1-0.5ug/ml
<b>Limitations</b>	This ACVRL1 antibody is available for research use only.




IHC staining of FFPE human gallbladder adenocarcinoma tissue with ACVRL1 antibody. HIER: boil tissue sections in pH8 EDTA for 20 min and allow to cool before testing.




IHC staining of FFPE human ovarian serous adenocarcinoma tissue with ACVRL1 antibody. HIER: boil tissue sections in pH8 EDTA for 20 min and allow to cool before testing.




IHC staining of FFPE human breast cancer tissue with ACVRL1 antibody. HIER: boil tissue sections in pH8 EDTA for 20 min and allow to cool before testing.



IHC staining of FFPE human liver cancer tissue with ACVRL1 antibody. HIER: boil tissue sections in pH8 EDTA for 20 min and allow to cool before testing.



Western blot testing of 1) human HL60, 2) human ThP-1, 3) rat brain, 4) rat heart, 5) mouse brain, 6) mouse kidney and 7) mouse heart tissue lysate with ACVRL1 antibody. Predicted molecular weight ~56 kDa.



Flow cytometry testing of human MCF7 cells with ACVRL1 antibody at 1ug/million cells (blocked with goat sera); Red=cells alone, Green=isotype control, Blue= ACVRL1 antibody.

## Description

Serine/threonine-protein kinase receptor R3 (SKR3) is an enzyme that in humans is encoded by the ACVRL1 gene. This gene encodes a type I cell-surface receptor for the TGF-beta superfamily of ligands. It shares with other type I receptors a high degree of similarity in serine-threonine kinase subdomains, a glycine- and serine-rich region (called the GS domain) preceding the kinase domain, and a short C-terminal tail. The encoded protein, sometimes termed ALK1, shares similar domain structures with other closely related ALK or activin receptor-like kinase proteins that form a subfamily of receptor serine/threonine kinases. Mutations in this gene are associated with hemorrhagic telangiectasia type 2, also known as Rendu-Osler-Weber syndrome 2.

## Application Notes

Optimal dilution of the ACVRL1 antibody should be determined by the researcher.

## Immunogen

Recombinant human protein (amino acids T271-L362) was used as the immunogen for the ACVRL1 antibody.

## Storage

After reconstitution, the ACVRL1 antibody can be stored for up to one month at 4oC. For long-term, aliquot and store at -20oC. Avoid repeated freezing and thawing.