


NDUFS8 Antibody / NADH-ubiquinone oxidoreductase 23 kDa subunit (RQ7236)


Catalog No.	Formulation	Size
RQ7236	0.5mg/ml if reconstituted with 0.2ml sterile DI water	100 ug

[Bulk quote request](#)


Availability	1-3 business days
Species Reactivity	Human, Mouse, Rat
Format	Antigen affinity purified
Host	Rabbit
Clonality	Polyclonal (rabbit origin)
Isotype	Rabbit IgG
Purity	Antigen affinity purified
Buffer	Lyophilized from 1X PBS with 2% Trehalose
UniProt	O00217
Localization	Cytoplasmic (mitochondria)
Applications	Western Blot : 0.5-1ug/ml Immunohistochemistry (FFPE) : 2-5ug/ml Flow Cytometry : 1-3ug/million cells Direct ELISA : 0.1-0.5ug/ml
Limitations	This NDUFS8 antibody is available for research use only.




Western blot testing of 1) rat heart, 2) rat liver, 3) rat skeletal muscle, 4) mouse heart, 5) mouse liver, 6) mouse skeletal muscle and 7) mouse C2C12 cell lysate with NDUFS8 antibody. Expected molecular weight ~23 kDa.




Western blot testing of human 1) HeLa, 2) HepG2, 3) A549, 4) MCF7, 5) Caco-2, 6) PC-3 and 7) 293T cell lysate with NDUF8 antibody. Expected molecular weight ~23 kDa.




IHC staining of FFPE human liver cancer tissue with NDUF8 antibody. HIER: boil tissue sections in pH8 EDTA for 20 min and allow to cool before testing.




IHC staining of FFPE mouse heart tissue with NDUF8 antibody. HIER: boil tissue sections in pH8 EDTA for 20 min and allow to cool before testing.



Immunofluorescent staining of FFPE human liver cancer tissue with NDUF8 antibody (red) and DAPI nuclear stain (blue). HIER: steam section in pH8 EDTA buffer for 20 min.



Flow cytometry testing of human 293T cells with NDUF8 antibody at 1ug/million cells (blocked with goat sera); Red=cells alone, Green=isotype control, Blue= NDUF8 antibody.



Flow cytometry testing of human HeLa cells with NDUF8 antibody at 1ug/million cells (blocked with goat sera); Red=cells alone, Green=isotype control, Blue= NDUF8 antibody.

Description

NADH dehydrogenase [ubiquinone] iron-sulfur protein 8, mitochondrial also known as NADH-ubiquinone oxidoreductase 23 kDa subunit, Complex I-23kD (CI-23kD), or TYKY subunit is an enzyme that in humans is encoded by the *NDUFS8* gene. This gene encodes a subunit of mitochondrial NADH:ubiquinone oxidoreductase, or Complex I, a multimeric enzyme of the respiratory chain responsible for NADH oxidation, ubiquinone reduction, and the ejection of protons from mitochondria. The encoded protein is involved in the binding of two of the six to eight iron-sulfur clusters of Complex I and, as such, is required in the electron transfer process. Mutations in this gene have been associated with Leigh syndrome.

Application Notes

Optimal dilution of the *NDUFS8* antibody should be determined by the researcher.

Immunogen

Recombinant human protein (amino acids M1-R210) was used as the immunogen for the *NDUFS8* antibody.

Storage

After reconstitution, the *NDUFS8* antibody can be stored for up to one month at 4°C. For long-term, aliquot and store at -20°C. Avoid repeated freezing and thawing.