


## ABI1 Antibody (F51187)


Catalog No.	Formulation	Size
F51187-0.4ML	In 1X PBS, pH 7.4, with 0.09% sodium azide	0.4 ml
F51187-0.08ML	In 1X PBS, pH 7.4, with 0.09% sodium azide	0.08 ml

[Bulk quote request](#)


<b>Availability</b>	1-3 business days
<b>Species Reactivity</b>	Human
<b>Predicted Reactivity</b>	Mouse, Rat
<b>Format</b>	Antigen affinity purified
<b>Host</b>	Rabbit
<b>Clonality</b>	Polyclonal (rabbit origin)
<b>Isotype</b>	Rabbit Ig
<b>Purity</b>	Antigen affinity
<b>UniProt</b>	Q8IZP0
<b>Localization</b>	Cytoplasmic
<b>Applications</b>	Western Blot : 1:1000 IHC (Paraffin) : 1:50-1:100 Flow Cytometry : 1:10-1:50 Immunofluorescence : 1:10-1:50
<b>Limitations</b>	This ABI1 antibody is available for research use only.




Western blot analysis of ABI1 antibody and human 1) MCF-7, 2) CEM, and 3) Jurkat lysate. Predicted molecular weight ~55 kDa, commonly observed at 55-65 kDa.



IHC analysis of FFPE human brain tissue stained with ABI1 antibody



Flow cytometric analysis of SK-Br-3 cells using ABI1 antibody (bottom histogram) compared to a [negative control](#) (top histogram). FITC-conjugated goat-anti-rabbit secondary Ab was used for the analysis.



Immunofluorescence analysis of ABI1 antibody with paraffin-embedded human brain tissue. Primary antibody was followed by FITC-conjugated goat anti-rabbit IgG (green).

## Description

Abelson interactor 1 has been found to form a complex with EPS8 and SOS1, and is thought to be involved in the transduction of signals from Ras to Rac. In addition, this protein may play a role in the regulation of EGF-induced Erk pathway activation as well as cytoskeletal reorganization and EGFR signaling.

## Application Notes

Titration of the ABI1 antibody may be required due to differences in protocols and secondary/substrate sensitivity.

## Immunogen

A portion of amino acids 81-108 from the human protein was used as the immunogen for this ABI1 antibody.

## Storage

Aliquot the ABI1 antibody and store frozen at -20oC or colder. Avoid repeated freeze-thaw cycles.