

ABO Antibody / Blood Group Antigen B [clone HEB-29] (V2551)

Catalog No.	Formulation	Size
V2551-100UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide	100 ug
V2551-20UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide	20 ug
V2551SAF-100UG	1 mg/ml in 1X PBS; BSA free, sodium azide free	100 ug



Citations (7)

[Bulk quote request](#)

Availability	1-3 business days
Species Reactivity	Human
Format	Purified
Clonality	Monoclonal (mouse origin)
Isotype	Mouse IgM, kappa
Clone Name	HEB-29
Purity	PEG precipitation
UniProt	P16442
Localization	Cell surface
Applications	Immunofluorescence : 2-4ug/ml Immunohistology (formalin-fixed) : 1-2ug/ml for 30 min at RT
Limitations	This ABO antibody is available for research use only.



Description

The antibody HEB-29 reacts with human blood group B. The specificity of the antibody HEB-29 was confirmed by

comparison of specificity and reactivity to standard reagent using >5.000 samples of blood. mAb HEB-29 shows specific staining of erythrocytes and vascular epithelium of blood group B controls and no staining in group A controls. It is applicable for tissue staining in tumor patients with blood groups B and AB. Blood group antigens are generally defined as molecules formed by sequential addition of saccharides to the carbohydrate side chains of lipids and proteins detected on erythrocytes and certain epithelial cells. The A, B and H antigens are reported to undergo modulation during malignant cellular transformation. Blood group related antigens represent a group of carbohydrate determinants carried on both glycolipids and glycoproteins. They are usually mucin type, and are detected on erythrocytes, certain epithelial cells, and in secretions of certain individuals. Sixteen genetically and biosynthetically distinct but inter related specificities belong to this group of antigens, including A, B, H, Lewis A, Lewis B, Lewis X, Lewis Y, and precursor type 1 chain antigens.

Application Notes

Immunogen

A mixture of erythrocytes of group B and glycoprotein fraction isolated from saliva of secretors with blood group B was used as the immunogen for the ABO antibody.

Storage

Store the ABO antibody at 2-8oC (with azide) or aliquot and store at -20oC or colder (without azide).