


## ABP1 Antibody / AOC1 / Amiloride binding protein 1 (R32508)


Catalog No.	Formulation	Size
R32508	0.5mg/ml if reconstituted with 0.2ml sterile DI water	100 ug

**Bulk quote request**


<b>Availability</b>	1-3 business days
<b>Species Reactivity</b>	Human, Monkey
<b>Format</b>	Antigen affinity purified
<b>Clonality</b>	Polyclonal (rabbit origin)
<b>Isotype</b>	Rabbit IgG
<b>Purity</b>	Antigen affinity
<b>Buffer</b>	Lyophilized from 1X PBS with 2% Trehalose
<b>UniProt</b>	P19801
<b>Localization</b>	Cytoplasmic, membranous
<b>Applications</b>	Western Blot : 0.5-1ug/ml Immunohistochemistry (FFPE) : 2-5ug/ml Immunofluorescence (FFPE) : 5ug/ml
<b>Limitations</b>	This ABP1 antibody is available for research use only.




IHC staining of FFPE human colonic adenoma tissue with ABP1 antibody. HIER: boil tissue sections in pH8 EDTA for 20 min and allow to cool before testing.




IHC staining of FFPE human placental tissue with ABP1 antibody. HIER: boil tissue sections in pH8 EDTA for 20 min and allow to cool before testing.



IHC staining of FFPE human renal cell carcinoma tissue with ABP1 antibody. HIER: boil tissue sections in pH8 EDTA for 20 min and allow to cool before testing.



Immunofluorescent staining of FFPE human T-47D cells with ABP1 antibody (green) and DAPI nuclear stain (blue). HIER: steam section in pH6 citrate buffer for 20 min.



Western blot testing of 1) human 293T, 2) human HK-2 and 3) monkey COS-7 lysate with ABP1 antibody. Predicted molecular weight ~85 kDa.

## Description

The AOC1 gene encodes a metal-binding membrane glycoprotein that oxidatively deaminates putrescine, histamine, and related compounds. The encoded protein is inhibited by amiloride, a diuretic that acts by closing epithelial sodium ion channels. Alternatively spliced transcript variants encoding multiple isoforms have been observed for this gene. Catalyzes the degradation of compounds such as putrescine, histamine, spermine, and spermidine, substances involved in allergic and immune responses, cell proliferation, tissue differentiation, tumor formation, and possibly apoptosis. Placental DAO is thought to play a role in the regulation of the female reproductive function.

## Application Notes

Differences in protocols and secondary/substrate sensitivity may require the ABP1 antibody to be titrated for optimal performance.

## Immunogen

Amino acids 144-180 (STAELYHTLQEATKPLHQFFLNTTGFSFQDCHDR) from the human protein were used as the immunogen for the ABP1 antibody.

## Storage

After reconstitution, the ABP1 antibody can be stored for up to one month at 4°C. For long-term, aliquot and store at -20°C. Avoid repeated freezing and thawing.

