


## ACADM Antibody / Mcad [Biotin Conjugate] (R35671BTN)

Catalog No.	Formulation	Size
R35671BTN	0.5 mg/ml in 1X TBS, pH7.3, with 0.5% BSA (US sourced) and 0.02% sodium azide	100 ug

[Bulk quote request](#)

<b>Availability</b>	1-3 business days
<b>Species Reactivity</b>	Human
<b>Format</b>	Biotin Conjugate
<b>Host</b>	Goat
<b>Clonality</b>	Polyclonal (goat origin)
<b>Isotype</b>	Goat Ig
<b>Purity</b>	Antigen affinity
<b>Gene ID</b>	34
<b>Applications</b>	Western Blot : 2-4ug/ml Immunohistochemistry (FFPE) : 3-5ug/ml ELISA (peptide) LOD : 1:32000
<b>Limitations</b>	This ACADM antibody is available for research use only.



Western blot testing of human heart lysate with ACADM antibody at 3ug/ml. Predicted molecular weight: ~46kDa. An NAP blocker was used in place of non-fat milk as blocking solution and diluent. (1)

### Description

Additional name(s) for this target protein: Acyl-Coenzyme A dehydrogenase medium chain

### Application Notes

Optimal dilution of the ACADM antibody should be determined by the researcher.

1. Do not use a blocking agent or diluent that contains biotin (including milk and some BSA preparations) when testing with biotinylated antibodies. A non-animal protein (NAP) blocker is recommended.

## **Immunogen**

Amino acids RLIVAREHIDKYKN were used as the immunogen for this ACADM antibody.

## **Storage**

Aliquot and store the ACADM antibody at -20oC.