


ACE2 Antibody / Angiotensin Converting Enzyme 2 [clone ACE2/7203] (V5342)


Catalog No.	Formulation	Size
V5342-100UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced), 0.05% sodium azide	100 ug
V5342-20UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced), 0.05% sodium azide	20 ug
V5342SAF-100UG	1 mg/ml in 1X PBS; BSA free, sodium azide free	100 ug

Bulk quote request

Availability	1-3 business days
Species Reactivity	Human
Format	Purified
Clonality	Monoclonal (mouse origin)
Isotype	Mouse IgG1, kappa
Clone Name	ACE2/7203
Purity	Protein A/G affinity
UniProt	Q9BYF1
Localization	Cell membrane, Secreted
Applications	ELISA (Order BSA-free Format For Coating) : Immunohistochemistry (FFPE) : 1-2ug/ml for 30 min at RT
Limitations	This ACE2 antibody is available for research use only.



IHC staining of FFPE human small intestine tissue with ACE2 antibody (clone ACE2/7203) at 2ug/ml. HIER: boil tissue sections in pH 9 10mM Tris with 1mM EDTA for 20 min and allow to cool before testing.



IHC staining of FFPE human kidney tissue with ACE2 antibody (clone ACE2/7203) at 2ug/ml. HIER: boil tissue sections in pH 9 10mM Tris with 1mM EDTA for 20 min and allow to cool before testing.

Description

Carboxypeptidase which converts angiotensin I to angiotensin 1-9, a peptide of unknown function, and angiotensin II to angiotensin 1-7, a vasodilator. Also able to hydrolyze apelin-13 and dynorphin-13 with high efficiency. By cleavage of angiotensin II, may be an important regulator of heart function. By cleavage of angiotensin II, may also have a protective role in acute lung injury. Plays an important role in amino acid transport by acting as binding partner of amino acid transporter SL6A19 in intestine, regulating trafficking, expression on the cell surface, and its catalytic activity. In case of human coronaviruses SARS and HCoV-NL63 infections, serve as functional receptor for the spike glycoprotein of both coronaviruses.

Application Notes

Optimal dilution of the ACE2 antibody should be determined by the researcher.

Immunogen

A recombinant partial protein sequence (within amino acids 1-300) from the human protein was used as the immunogen for the ACE2 antibody.

Storage

Aliquot the ACE2 antibody and store frozen at -20°C or colder. Avoid repeated freeze-thaw cycles.