


## Acidic Cytokeratin Antibody [clone ACCK1-1] (V7193)


Catalog No.	Formulation	Size
V7193-100UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide	100 ug
V7193-20UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide	20 ug
V7193SAF-100UG	1 mg/ml in 1X PBS; BSA free, sodium azide free	100 ug
V7193IHC-7ML	Prediluted in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide; *For IHC use only*	7 ml

[Bulk quote request](#)

<b>Availability</b>	1-3 business days
<b>Species Reactivity</b>	Human, Rat
<b>Format</b>	Purified
<b>Host</b>	Mouse
<b>Clonality</b>	Monoclonal (mouse origin)
<b>Isotype</b>	Mouse IgG1, kappa
<b>Clone Name</b>	ACCK1-1
<b>Purity</b>	Protein G affinity chromatography
<b>UniProt</b>	Q7Z794
<b>Localization</b>	Cytoplasmic
<b>Applications</b>	Immunohistochemistry (FFPE) : 1-2ug/ml for 30 min at RT (1) Prediluted IHC Only Format : incubate for 30 min at RT (2)
<b>Limitations</b>	This Acidic Cytokeratin antibody is available for research use only.



IHC testing of FFPE human tonsil with Acidic Cytokeratin antibody.



IHC testing of FFPE rat stomach with Acidic Cytokeratin antibody.

## Description

There are two types of cytokeratins/keratins/CKs: the acidic type I cytokeratins and the basic or neutral type II cytokeratins. The subsets of cytokeratins which an epithelial cell expresses depends mainly on the type of epithelium, the moment in the course of terminal differentiation and the stage of development. Thus this specific keratin fingerprint allows the classification of all epithelia upon their keratin expression profile. Furthermore this applies also to the malignant counterparts of the epithelia (carcinomas), as the keratin profile tends to remain constant when an epithelium undergoes malignant transformation. The main clinical implication is that the study of the keratin profile by immunohistochemistry techniques is a tool of immense value widely used for tumor diagnosis and characterization in surgical pathology. [Wiki]

## Application Notes

Titering of the Acidic Cytokeratin antibody may be required for optimal performance.

1. FFPE testing requires sections to be boiled in pH6 10mM citrate buffer for 10-20 minutes, followed by cooling at RT for 20 minutes, prior to staining.
2. The prediluted format is supplied in a dropper bottle and is optimized for use in IHC. After epitope retrieval step (if required), drip mAb solution onto the tissue section and incubate at RT for 30 min.
3. This antibody will detect the following cytokeratins (CK): CK10: 56kDa; CK14: 50kDa; CK15: 50kDa; CK16: 48kDa; CK19: 40kDa.

## Immunogen

An amino acid sequence common to acidic/type I keratins was used as the immunogen for the Acidic Cytokeratin antibody.

## Storage

Store the Acidic Cytokeratin antibody at 2-8oC (with azide) or aliquot and store at -20oC or colder (without azide).