


Adam28 Antibody (RQ6034)


Catalog No.	Formulation	Size
RQ6034	0.5mg/ml if reconstituted with 0.2ml sterile DI water	100 ug

Bulk quote request


Availability	1-3 business days
Species Reactivity	Mouse, Rat
Format	Antigen affinity purified
Host	Rabbit
Clonality	Polyclonal (rabbit origin)
Isotype	Rabbit IgG
Purity	Affinity purified
Buffer	Lyophilized from 1X PBS with 2% Trehalose and 0.025% sodium azide
UniProt	E9PTQ3
Applications	Western Blot : 0.5-1ug/ml Immunohistochemistry : 1-2ug/ml Flow Cytometry : 1-3ug/million cells Direct ELISA : 0.1-0.5ug/ml
Limitations	This Adam28 antibody is available for research use only.



IHC staining of FFPE rat gastric tissue with Adam28 antibody. HIER: boil tissue sections in pH8 EDTA for 20 min and allow to cool before testing.




IHC staining of FFPE mouse lung tissue with Adam28 antibody. HIER: boil tissue sections in pH8 EDTA for 20 min and allow to cool before testing.



IHC staining of FFPE mouse lung tissue with Adam28 antibody. HIER: boil tissue sections in pH8 EDTA for 20 min and allow to cool before testing.

KDa
180-
130-
95-
72-
55-
43-
34-
26-
17-

Western blot testing of rat brain lysate with Adam28 antibody. Predicted molecular weight ~87 kDa.



Flow cytometry testing of mouse HEPA1-6 cells with Adam28 antibody at 1ug/million cells (blocked with goat sera); Red=cells alone, Green=isotype control, Blue= Adam28 antibody.

Description

Disintegrin and metalloproteinase domain-containing protein 28 is an enzyme that in humans is encoded by the ADAM28 gene. This gene encodes a member of the ADAM (a disintegrin and metalloprotease domain) family. Members of this family are membrane-anchored proteins structurally related to snake venom disintegrins, and have been implicated in a variety of biological processes involving cell-cell and cell-matrix interactions, including fertilization, muscle development, and neurogenesis. The protein encoded by this gene is a lymphocyte-expressed ADAM protein. And this gene is present in a gene cluster with other members of the ADAM family on chromosome 8. Alternative splicing results in multiple transcript variants.

Application Notes

Optimal dilution of the Adam28 antibody should be determined by the researcher.

Immunogen

Recombinant rat protein (amino acids K26-K195) was used as the immunogen for the Adam28 antibody.

Storage

After reconstitution, the Adam28 antibody can be stored for up to one month at 4°C. For long-term, aliquot and store at -20°C. Avoid repeated freezing and thawing.