


## ADH1C Antibody / Alcohol dehydrogenase 1C (F54417)


Catalog No.	Formulation	Size
F54417-0.4ML	In 1X PBS, pH 7.4, with 0.09% sodium azide	0.4 ml
F54417-0.08ML	In 1X PBS, pH 7.4, with 0.09% sodium azide	0.08 ml

**Bulk quote request**


<b>Availability</b>	1-3 business days
<b>Species Reactivity</b>	Human
<b>Format</b>	Purified
<b>Clonality</b>	Polyclonal (rabbit origin)
<b>Isotype</b>	Rabbit Ig
<b>Purity</b>	Antigen affinity purified
<b>UniProt</b>	P00326
<b>Localization</b>	Cytoplasmic
<b>Applications</b>	Flow Cytometry : 1:25 (1x10e6 cells) Immunofluorescence : 1:25 Immunohistochemistry (FFPE) : 1:25 Western Blot : 1:500-1:2000
<b>Limitations</b>	This ADH1C antibody is available for research use only.




Western blot testing of human T-47D cell lysate with ADH1C antibody. Predicted molecular weight ~40 kDa.




Western blot testing of 1) non-transfected and 2) transfected 293 cell lysate with ADH1C antibody.



Immunofluorescent staining of human T-47D cells with ADH1C antibody (green) and DAPI nuclear stain (blue).



IHC testing of FFPE human hepatocarcinoma tissue with ADH1C antibody. HIER: steam section in pH6 citrate buffer for 20 min and allow to cool prior to staining.



Flow cytometry testing of human CCRF-CEM cells with ADH1C antibody; Blue=isotype control, Green= ADH1C antibody.

## Description

This gene encodes class I alcohol dehydrogenase, gamma subunit, which is a member of the alcohol dehydrogenase family. Members of this enzyme family metabolize a wide variety of substrates, including ethanol, retinol, other aliphatic alcohols, hydroxysteroids, and lipid peroxidation products. Class I alcohol dehydrogenase, consisting of several homo- and heterodimers of alpha, beta, and gamma subunits, exhibits high activity for ethanol oxidation and plays a major role in ethanol catabolism. Three genes encoding alpha, beta and gamma subunits are tandemly organized in a genomic segment as a gene cluster.

## Application Notes

The stated application concentrations are suggested starting points. Titration of the ADH1C antibody may be required due to differences in protocols and secondary/substrate sensitivity.

## Immunogen

A portion of amino acids 231-260 from the human protein was used as the immunogen for the ADH1C antibody.

## Storage

Aliquot the ADH1C antibody and store frozen at -20oC or colder. Avoid repeated freeze-thaw cycles.