


# AFP Antibody Cocktail [clone C2 + C3 + MBS-12] (V2476)

Catalog No.	Formulation	Size
V2476-100UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide	100 ug
V2476-20UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide	20 ug
V2476SAF-100UG	1 mg/ml in 1X PBS; BSA free, sodium azide free	100 ug

## **Bulk quote request**

Availability	1-3 business days
Species Reactivity	Human
Format	Purified
Clonality	Monoclonal (mouse origin)
Isotype	Mouse IgG1, kappa
Clone Name	C2 + C3 + MBS-12
Purity	Protein G affinity chromatography
UniProt	P02771
Localization	Cytoplasmic
Applications	Immunofluorescence : 1-2ug/ml Immunohistochemistry (FFPE) : 1-2ug/ml for 30 min at RT
Limitations	This AFP antibody cocktail is available for research use only.



IHC: Formalin-fixed, paraffin-embedded human fetal liver stained with AFP antibody (C2  $\pm$  C3  $\pm$  MBS-12).

## **Description**

mAb is highly specific to AFP and shows no cross-reaction with other oncofetal antigens or serum albumin. AFP is normally synthesized in the liver, intestinal tract, and yolk sac of the fetus. Antibody to AFP has been shown to be useful in detecting hepatocellular carcinomas (HCC) and germ cell neoplasms, especially yolk sac tumors.

### **Application Notes**

Optimal dilution of the AFP antibody cocktail should be determined by the researcher.

- 1. Staining of formalin-fixed tissues requires boiling tissue sections in 10mM Citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 min
- 2. The prediluted format is supplied in a dropper bottle and is optimized for use in IHC. After epitope retrieval step (if required), drip mAb solution onto the tissue section and incubate at RT for 30 min.

#### **Immunogen**

Recombinant full-length human Alpha Fetoprotein was used as the immunogen for the AFP antibody cocktail.

#### **Storage**

Store the AFP antibody cocktail at 2-8oC (with azide) or aliquot and store at -20oC or colder (without azide).