


AGER Antibody / RAGE [clone 5C6C1] (RQ7676)


Catalog No.	Formulation	Size
RQ7676	0.5mg/ml if reconstituted with 0.2ml sterile DI water	100 ug

Bulk quote request


Availability	1-3 business days
Species Reactivity	Human, Mouse, Rat
Format	Antigen affinity purified
Host	Mouse
Clonality	Monoclonal (mouse origin)
Isotype	Mouse IgG2b
Clone Name	5C6C1
Purity	Antigen affinity purified
Buffer	Lyophilized from 1X PBS with 2% Trehalose
UniProt	Q15109
Applications	Western Blot : 0.5-1ug/ml Immunohistochemistry (FFPE) : 2-5ug/ml Flow Cytometry : 1-3ug/million cells
Limitations	This AGER antibody is available for research use only.




IHC staining of FFPE mouse lung tissue with AGER antibody. HIER: boil tissue sections in pH8 EDTA for 20 min and allow to cool before testing.




IHC staining of FFPE rat lung tissue with AGER antibody. HIER: boil tissue sections in pH8 EDTA for 20 min and allow to cool before testing.



IHC staining of FFPE rat lung tissue with AGER antibody. HIER: boil tissue sections in pH8 EDTA for 20 min and allow to cool before testing.



Western blot testing of 1) rat lung and 2) mouse lung tissue lysate with AGER antibody. Predicted molecular weight ~43 kDa.



Flow cytometry testing of human Jurkat cells with AGER antibody at 1ug/million cells (blocked with goat sera); Red=cells alone, Green=isotype control, Blue= AGER antibody.

Description

The receptor for advanced glycation end products (RAGE) is a multi-ligand member of the immunoglobulin superfamily of cell surface molecules. It interacts with distinct molecules implicated in homeostasis, development and inflammation, and certain diseases such as diabetes and Alzheimer's disease. RAGE is also a central cell surface receptor for amphotericin and EN-RAGE. And RAGE is associated with sustained NF-kappaB activation in the diabetic microenvironment and has a central role in sensory neuronal dysfunction. Moreover, RAGE propagates cellular dysfunction in several inflammatory disorders and diabetes, and it also functions as an endothelial adhesion receptor promoting leukocyte recruitment.

Application Notes

Optimal dilution of the AGER antibody should be determined by the researcher.

Immunogen

Amino acids 91-120 (IQDEGIFRCQAMNRNGKETKSNYRVRVYQI) from the human protein were used as the immunogen for the AGER antibody.

Storage

After reconstitution, the AGER antibody can be stored for up to one month at 4oC. For long-term, aliquot and store at -20oC. Avoid repeated freezing and thawing.