


Albumin Antibody / ALB (F54330)


Catalog No.	Formulation	Size
F54330-0.2ML	In 1X PBS, pH 7.4, with 0.09% sodium azide	0.2 ml
F54330-0.05ML	In 1X PBS, pH 7.4, with 0.09% sodium azide	0.05 ml

[Bulk quote request](#)


Availability	1-3 business days
Species Reactivity	Human, Mouse
Format	Purified
Host	Rabbit
Clonality	Polyclonal (rabbit origin)
Isotype	Rabbit Ig
Purity	Antigen affinity purified
UniProt	P02768
Applications	Immunohistochemistry (FFPE) : 1:25 Western Blot : 1:500-1:2000 Flow Cytometry : 1:25 (1x10 ⁶ cells)
Limitations	This Albumin antibody is available for research use only.




Western blot testing of 1) human HepG2, 2) human HeLa and 3) mouse liver lysate with Albumin antibody. Predicted molecular weight ~66 kDa.



Albumin Antibody Human Kidney Tissue IHC. Immunohistochemistry staining of FFPE human kidney tissue with Albumin antibody. HIER: steam section in pH9 EDTA for 20 min and allow to cool prior to staining.



Albumin Antibody Human Liver Tissue IHC. Immunohistochemistry staining testing of FFPE human liver tissue with Albumin antibody. HIER: steam section in pH9 EDTA for 20 min and allow to cool prior to staining.



Albumin Antibody HepG2 FACS. Flow cytometry testing of human HepG2 cells with Albumin antibody; Blue=isotype control, Green= Albumin antibody.

Description

Albumin Antibody detects Albumin, a soluble, monomeric protein which comprises about one-half of the blood serum protein. Albumin functions primarily as a carrier protein for steroids, fatty acids, and thyroid hormones and plays a role in stabilizing extracellular fluid volume. Albumin is a globular unglycosylated serum protein of molecular weight 65,000. Albumin is synthesized in the liver as preproalbumin which has an N-terminal peptide that is removed before the nascent protein is released from the rough endoplasmic reticulum. The product, proalbumin, is in turn cleaved in the Golgi vesicles to produce the secreted albumin.

For broader detection of Albumin in liver function and hepatocyte biology studies, see our [Albumin antibody \(clone ALB/2141\)](#).

Application Notes

The stated application concentrations are suggested starting points. Titration of the Albumin antibody may be required due to differences in protocols and secondary/substrate sensitivity.

Immunogen

A portion of amino acids 540-569 from the human protein was used as the immunogen for the Albumin antibody.

Storage

Aliquot the Albumin antibody and store frozen at -20°C or colder. Avoid repeated freeze-thaw cycles.